

Research



Cite this article: Gaos AR *et al.* 2017 Natal foraging philopatry in eastern Pacific hawksbill turtles. *R. Soc. open sci.* 4: 170153. <http://dx.doi.org/10.1098/rsos.170153>

Received: 22 February 2017

Accepted: 26 July 2017

Subject Category:

Genetics

Subject Areas:

ecology/developmental biology/genetics

Keywords:

natal homing, life history, hatchling dispersal, spatial ecology, conservation genetics, juvenile

Author for correspondence:

Alexander R. Gaos

e-mail: gaos@hawksbill.org

[†]Present address: 4314 Dalles Court, San Diego, CA 92117, USA.

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.3852109>.

Natal foraging philopatry in eastern Pacific hawksbill turtles

Alexander R. Gaos^{1,2,3,26,†}, Rebecca L. Lewison¹, Michael P. Jensen³, Michael J. Liles^{4,5,26}, Ana Henriquez^{5,26}, Sofia Chavarria^{5,26}, Carlos Mario Pacheco^{5,26}, Melissa Valle^{5,26}, David Melero²⁶, Velkiss Gadea^{6,26}, Eduardo Altamirano^{6,26}, Perla Torres^{7,26}, Felipe Vallejo^{8,26}, Cristina Miranda^{8,26}, Carolina LeMarie^{8,26}, Jesus Lucero^{9,26}, Karen Ocegüera^{9,26}, Didiher Chácon^{10,26}, Luis Fonseca^{10,26}, Marino Abrego^{11,26}, Jeffrey A. Seminoff^{12,26}, Eric E. Flores^{13,14}, Israel Llamas^{15,26}, Rodrigo Donadi¹⁶, Bernardo Peña¹¹, Juan Pablo Muñoz^{17,26}, Daniela Alarcón Ruales^{17,26}, Jaime A. Chaves¹⁷, Sarah Otterstrom^{18,26}, Alan Zavala^{19,20,26}, Catherine E. Hart^{21,26}, Rachel Brittain^{22,26}, Joanna Alfaro-Shigueto^{23,24,25,26}, Jeffrey Mangel^{23,26}, Ingrid L. Yañez²⁶ and Peter H. Dutton¹²

¹Department of Biology, San Diego State University, San Diego, CA, USA

²Graduate Group in Ecology, University of California Davis, Davis, CA, USA

³Marine Mammal and Turtle Division, Ocean Associates Inc., under contract to the Southwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, La Jolla, CA, USA

⁴Department of Biology, University of Texas at El Paso, El Paso, TX, USA

⁵ProCosta, San Salvador, El Salvador

⁶Marine Turtles Department, Fauna & Flora International, Managua, Nicaragua

⁷Instituto de Ciencias del Mar y Limnología, Unidad Académica Mazatlán, Universidad Nacional de México, Mazatlán, México


⁸Equilibrio Azul, Quito, Ecuador

⁹Grupo Tortuguero de las Californias, A.C., La Paz, México

¹⁰Latin American Sea Turtles, Tibás, Costa Rica

¹¹Conservación de Recursos Costeros y Marinos, Ministerio del Ambiente de Panamá, Panama City, Panama

- ¹²Marine Mammal and Turtle Division, Southwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, La Jolla, CA, USA
- ¹³Sistema Nacional de Investigación, Panama City, Panama
- ¹⁴Instituto de Investigaciones Científicas y Servicios de Alta Tecnología, Panama City, Panama
- ¹⁵Campamento Tortuguero Mayto, A.C., Mayto, Mexico
- ¹⁶World Wildlife Fund of Panama, Panama City, Panama
- ¹⁷Marine Ecology Department, Universidad San Francisco de Quito/Galapagos Science Center, San Cristóbal, Galapagos Archipelago, Ecuador
- ¹⁸Paso Pacífico, Managua, Nicaragua
- ¹⁹Unidad Sinaloa, Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional, Sinaloa, Mexico
- ²⁰Instituto Politécnico Nacional, Sinaloa, Mexico
- ²¹Red Tortuguera, A.C, Guayabitos, Mexico
- ²²Akazul, La Barrona, Guatemala
- ²³Marine Turtle Research Group, School of Biosciences, University of Exeter, Penryn, UK
- ²⁴Marine Biology Department, Universidad Científica del Sur, Lima, Peru
- ²⁵ProDelphinus, Lima, Peru
- ²⁶Eastern Pacific Hawksbill Initiative, San Diego, CA, USA

 ARG, 0000-0001-6100-7319

The complex processes involved with animal migration have long been a subject of biological interest, and broad-scale movement patterns of many marine turtle populations still remain unresolved. While it is widely accepted that once marine turtles reach sexual maturity they home to natal areas for nesting or reproduction, the role of philopatry to natal areas during other life stages has received less scrutiny, despite widespread evidence across the taxa. Here we report on genetic research that indicates that juvenile hawksbill turtles (*Eretmochelys imbricata*) in the eastern Pacific Ocean use foraging grounds in the region of their natal beaches, a pattern we term natal foraging philopatry. Our findings confirm that traditional views of natal homing solely for reproduction are incomplete and that many marine turtle species exhibit philopatry to natal areas to forage. Our results have important implications for life-history research and conservation of marine turtles and may extend to other wide-ranging marine vertebrates that demonstrate natal philopatry.

1. Introduction

Understanding movement patterns is an essential component of successful wildlife management [1,2]. The movement and dispersal of highly migratory marine organisms such as marine turtles have been studied for decades (e.g. [3,4]) and some broad-scale patterns have become widely accepted. For instance, it is generally believed that most post-hatchling marine turtles spend a multi-year period in open-ocean gyres before recruiting to neritic foraging grounds, but deviations from this pattern are known to exist for some species and populations [5–9]. An abundance of research has also shown that once reaching sexual maturity, which can take between 10 and 50 years [9], most turtles return to the vicinity of their natal beaches to nest or reproduce (i.e. natal homing *sensu* [10]).

The specific dispersal behaviour of marine turtles to foraging grounds in relation to natal nesting beaches, termed rookeries, is not defined [11]. Particle simulation modelling and satellite tracking of post-hatchling turtles suggest a range of dispersal pathways over thousands of kilometres, indicating settlement in foraging areas can occur over extremely large spatial scales (e.g. [12–15]). Genetic research has demonstrated that foraging grounds are typically made up of individuals from multiple rookeries, further highlighting the potentially diffuse distribution of post-hatchling turtles [11]. Recent studies have found that juvenile loggerhead (*Caretta caretta*), green (*Chelonia mydas*) and hawksbill (*Eretmochelys imbricata*) turtles in the Atlantic actively home to foraging grounds in the vicinity of their natal habitats [16–22]. Despite evidence supporting this homing pattern across several marine turtle species, current literature on general marine turtle life-history has not fully recognized philopatry to natal areas during early life stages or explored potential conservation implications of this pattern [9,23].

In this study, we analysed mitochondrial DNA (mtDNA) sequences of juvenile and adult hawksbills from the eastern Pacific Ocean to evaluate the species' distribution at foraging grounds relative to natal rookeries and to assess the evidence of foraging philopatry to natal areas. Hawksbill turtles were considered virtually absent from the eastern Pacific Ocean as recently as 2007 [24] and remain as one of




Figure 1. Sample collection sites, including source rookeries (grey triangles), foraging grounds with sample sizes in the range $n = 20$ – 117 (white stars) that were used in principal components analysis to determine regional foraging grounds (RFGs; black boxes), as well as foraging grounds with sample sizes in the range $n = 1$ – 7 (black circles) that were also included in each RFG.

the world's most threatened marine turtle regional management units [25]. Although extensive field research over the past 10 years has revealed important hawksbill rookeries and foraging grounds, there are still considerable gaps in our understanding of dispersal of post-hatchling turtles to foraging areas and how rookeries contribute to foraging grounds. Additionally, several unique life-history characteristics have been discovered for hawksbills in this ocean region that could potentially influence the dispersal of hatchlings and rookery (i.e. nesting colony) contributions, including highly neritic habitat use, restricted home ranges and limited migrations [26–28], heightening interest in genetic research of hawksbills at foraging grounds.

2. Material and methods

2.1. Sample collection and processing

We collected tissue samples from hawksbill turtles at foraging grounds along the eastern Pacific rim (figure 1) between 2004 and 2015 using a combination of direct monitoring (i.e. tangle nets or manual dive captures), strandings and fisheries by-catch monitoring activities. We used rookery samples from Gaos *et al.* [29], which were supplemented by seven additional rookery samples. When feasible, we measured the curved carapace length (CCL) for all hawksbills encountered and applied Inconel flipper tags (National Band and Tag, Newport, KY, USA) to allow for ongoing identification. After collection, samples were placed in vials containing greater than 95% ethanol or water saturated with sodium chloride, which were subsequently stored in a -20°C freezer.

We amplified a 766bp segment of mtDNA from the control region of hawksbill tissue samples following the protocol outlined in Gaos *et al.* [29]. Haplotypes (KT934080, KT934070, KT964296,




Figure 2. Curved carapace length (CCL) distribution bins (centimetres) of hawksbills analysed in this study. Asterisk indicates bin containing average CCL for nesting female hawksbills in the eastern Pacific [34].

KR012503, KR012504, KT003685, KT072795, KR012505, KT072797, KU695258, KU695259, KX646708, KT934051) were determined using GENEIOUS v. R8 (Biomatters Inc., Newark, NJ, USA) and aligned against sequences in a local reference library, as well as those found in the GenBank database (<http://www.ncbi.nlm.nih.gov>).

2.2. Data analysis

We calculated F_{ST} values for rookeries and foraging grounds using ARLEQUIN v. 3.5.1.2 [30]. Because we were interested in identifying broad-scale genetic patterns and had variable sample sizes (mean $n = 12.4$, s.d. = 24.8) from disparate foraging grounds, we pooled samples to create regional foraging grounds (RFGs). To determine which samples would be included in each RFG, we conducted a principal components analysis (PCA) in the package GENALEX v. 6.5 [31] on F_{ST} values for eight foraging grounds for which we had sample sizes in the range $n = 20$ –117 (figure 1). The samples from the foraging grounds in each PCA cluster were then combined with neighbouring foraging grounds for which we had sample sizes in the range $n = 1$ –7 (figure 1).

We used Bayesian mixed stock analysis (MSA) in the software ‘BAYES’ [32] to estimate the contribution of rookeries to each RFG. The two rookeries in RFG2 were combined into a single regional estimate due to both their proximity and genetic similarity. Average annual number of nests at rookeries in RFG1 = 6.9 (± 7.3), RFG2 = 187.6 (± 50.7), RFG3 = 52.0 ($\pm n.a.$) and RFG4 = 22.3 (± 10.0) [33]. For the MSA, we ran a total of four Markov chain Monte Carlo (MCMCs) using uniform priors, coinciding with the number of potential source rookeries, with each MCMC consisting of 50 000 steps initiated at different starting points. We used a burn-in of 25 000 steps and calculated the posterior distribution from the remaining 25 000 for all chains, then ran the Gelman and Rubin shrink factor diagnostic to test that all chains had converged (i.e. less than 1.2) [32]. Individuals with haplotypes not observed in rookeries (i.e. orphan haplotypes, $n = 5$, 0.9% of total samples) were removed from the study.

3. Results

We obtained 766 bp sequences from the mtDNA control region for hawksbill turtles samples collected at 45 foraging grounds ($n = 535$) and five rookeries ($n = 267$) in nine countries (figure 1). Samples collected at foraging grounds included hawksbills with average CCL 51.4 cm (s.d. = 14.1, range 14.0–95.0, figure 2).

Our PCA identified four distinct geographical clusters (see electronic supplementary material, figure 1 for PCA results), leading to the creation of four RFGs (mean sample size $n = 133.8$, s.d. = 78.7). F_{ST}




Figure 3. Results of the Bayesian mixed stock analysis showing the contributions of rookeries based on location (represented by bar colour) to the four regional foraging grounds (RFGs). Nearest rookeries (blue bars) contributed the majority of foraging turtles to the RFG in which they were found.

values indicated highly significant ($p < 0.001$) structure among all potential source rookeries except for the two located in RFG1 and RFG4. However, these rookeries are separated by greater than 3500 km and it is extremely unlikely that they are demographically homogeneous [35], thus both were treated as independent source rookeries in our MSA.

Results of our MSA indicated that rookeries located in each RFG (figure 1) contribute the overwhelming majority (average = 80.5%, s.d. = 10.5) of turtles to foraging grounds in that same RFG (figure 3; see electronic supplementary material, table S1 for detailed MSA results).

4. Discussion

Although it has long been established that upon reaching sexual maturity, marine turtles return to the vicinity of their natal beaches to nest or reproduce [36], the potential role of philopatry to foraging areas during juvenile stages has remained largely overlooked as a general life-history trait within the taxon. Our results indicate that juvenile hawksbills in the eastern Pacific use foraging grounds in the region of their source rookeries, a pattern we term natal foraging philopatry (NFP). Despite major differences in the size of our source rookeries, i.e. annual nests among rookeries varied in some cases by an order of magnitude, our results indicate that rookeries in each region, irrespective of their size, were the primary contributors to foraging grounds in the same regions in which they were found. Similarly, while our results indicate rookery contributions are influenced by distance, our findings demonstrate that rookeries in each region contributed virtually all turtles to foraging grounds in those same regions (figure 3). This finding was consistent even for RFGs that were in close proximity to one another (e.g. RFG2 and RFG3).

Our results confirm recent studies that found evidence of NFP in Atlantic juvenile loggerhead, green and hawksbill turtles [16–22]. Combined, these findings suggest that natal philopatry for marine turtles

extends beyond mating and nesting adults, and that NFP is a common strategy exhibited across life stages for a number of marine turtle species and stocks. NFP has not been described in all species, e.g. leatherback sea turtles (*Dermochelys coriacea*), yet the pervasiveness of the NFP pattern is particularly notable given that it can manifest even in species and populations known to experience a panmictic post-hatchling dispersal pattern (e.g. [16,20,37]).

Although the NFP pattern is clearly observed in eastern Pacific hawksbills, what is not clear is the role of potential mechanisms behind the NFP pattern in this and other species or stocks, which could include oceanic currents, geomagnetic imprinting, visual and chemical cues, genetic memory, habitat associations, learned migration behaviour and other factors [9,15,38–40]. Our findings of NFP and previous field research [26,27,29,33,41] suggest that eastern Pacific hawksbills may not conform to the commonly postulated open-ocean dispersal theory in which post-hatchling turtles are believed to leave natal areas via entrainment in offshore currents [7,9,37], which for hawksbills would be major ocean currents of the eastern Pacific [42]. Instead, we raise the possibility that hawksbill in the eastern Pacific remain in coastal regions during the post-hatchling phase, categorized by Bolten [9] as a Type 1 hatchling developmental pattern, or undergo a truncated pelagic dispersal phase that keeps them out of major offshore currents. In contrast with marine turtle populations in other parts of the world, hawksbills along Pacific Central America nest at rookeries located within inshore mangrove estuaries [43,44] and these systems are strongly regulated by tidal fluctuations. Coupled with the reduced swim gaits exhibited by hawksbills compared with other marine turtle species [45], these currents could maintain hawksbill hatchlings closer to shore during the initial post-hatching phase in the eastern Pacific [46]. Recent research of lobster gillnet fisheries that operate in neritic waters (i.e. less than 1 km from shore) off Pacific Central America has led to the documentation of various hawksbills with CCL measuring 15 cm or less in size [41], further supporting the Type 1 hatchling development model as a potential mechanism underlying NFP for hawksbills in the eastern Pacific. To date, the flatback turtle (*Natator depressus*) is the only species of marine turtle whose hatchlings have been identified as not having a pelagic phase during early post-hatchling development [9,41,47], a pattern that has been largely attributed to the inshore location of their nesting beaches [46]. At present, data on fine-scale currents are not available for the coastal zones of the eastern Pacific where hawksbills are found, which limits opportunities to conduct meaningful particle modelling efforts. Future research combining particle modelling (e.g. [12,14]), tracking of post-hatchlings (e.g. [34]) and satellite telemetry of young life-stage (6–12 months old) juveniles (e.g. [15,37]) would help elucidate the post-hatchling movements of hawksbills in the eastern Pacific.

Irrespective of the mechanism, affinity to foraging grounds in proximity to rookeries would provide several selective advantages, namely higher lifetime reproductive success from more frequent visits to rookeries [48], as well as higher survival as the absence of large-scale migrations between rookeries and foraging grounds would reduce some threat exposure [4,14,49]. Recent telemetry and nesting biology research of female hawksbills in the eastern Pacific confirms relatively short post-nesting migrations or non-migratory behaviour altogether [26], as well as relatively common annual nesting intervals [29].

Our findings have important implications for life-history research and conservation of marine turtles, particularly for hawksbills in the eastern Pacific. NFP may mean that addressing local threats to foraging hawksbills has potentially greater conservation value for nearby rookeries and hence local populations. Similarly, whereas the conservation of marine turtles can often require the collaboration of governments from multiple nations over large ocean regions [49–51], NFP of hawksbills in the eastern Pacific indicates that one or two neighbouring nations may be able to protect the majority of the life cycle of an entire stock or management unit. Additionally, large rookeries are often recognized for their importance to sustaining marine turtle populations [52] by provisioning a ‘storage effect’, in which via one or two solid recruitment classes per generation they can enable repopulation of an ocean region on ecological timescales [29,53]. However, findings of NFP and the strong stock structure among RFGs indicate that rookeries in each RFG maintain unique genetic contributions, which suggests that focusing conservation efforts solely on larger rookeries would be misguided.

As NFP appears common among many marine turtle species, focused research on the ubiquity of this pattern in individual populations is needed to advance marine turtle ecology and to support effective conservation management strategies. Similarly, NFP as a general life-history concept merits specific discussion and inclusion in current marine turtle and spatial ecology literature. Finally, further research into NFP for other wide-ranging aquatic species known to exhibit nesting philopatry, such as

various salmonid and seabird species (e.g. [54,55]), may also be warranted to support robust conservation planning.

Ethics. All hawksbill handling procedures were approved (APF no.: 09-05-015 L—San Diego State University) by the Institutional Animal Care and Use Committee (IACUC) in accordance with the requirements pertaining to animal subjects protections within the Public Health Service Policy and USDA Animal Welfare Regulations. Scientific permits to collect hawksbill tissue samples were obtained in all countries and include the following permit numbers: Mexico (SGPA/DGVS: 03540/08, 03650/09, 04568/11, 05137/12, 02259/14, 04478/15, 07584/16), Guatemala (004/2014), El Salvador (MARN: AIMA-38-2008, DGOA-GE-089-2012, DGBPN-GVS-016-2013, DEV-GVS-17-2015), Nicaragua (003-012010, 008-042011, 003-012012, 006-062013, 001-012014, 006042015, 009-2016), Costa Rica (481-2012, ACG-PI-012-2014), Panama (SE/A-77-14, SE/A-17-15), Ecuador (011-DRM-MA, MAE-CGZ4-DPAM-2014-3286, MAE-DPASE-2016-0152, PC-35-12) and Peru (068-2004-INRENA-IFFS-DCB). Tissue import (United States (US844694/9)) and export (sample collection country) permits to meet requirements established by CITES were acquired from the appropriate environmental authorities in each country, including CONANP in Mexico (MX 75768), MARN in Guatemala (000653), MARN in El Salvador (09977), MARENA in Nicaragua (11357), MINAET in Costa Rica (2015-CR1764/SJ (#S 1824)), ANAM in Panama (SEXA-06-11), MAE in Ecuador (106/BG) and MINAM in Peru (6275). All research participants received consent from the aforementioned environmental authorities in their respective countries.

Data accessibility. The input files used in the MSA are available via the Dryad repository at: (<http://dx.doi.org/10.5061/dryad.38t23>) [56]. The sequence files used in this analysis (GenBank accession numbers: KT934070, KT964296, KR012503, KR012504, KT003685, KT072795, KR012505, KT072797, KU695258, KU695259, KX646708, KT934051) are available through the GenBank database (<http://www.ncbi.nlm.nih.gov>).

Authors' contributions. A.R.G. carried out laboratory work and drafted the manuscript; R.L.L., P.H.D., M.J.L., M.P.J., J.A.S., J.A.-S. and J.M. helped develop the manuscript; A.R.G. and M.P.J. conducted the analyses; A.R.G., M.J.L., A.H., S.C., C.M.P., M.V., D.M., V.G., E.A., P.T., F.V., C.M., C.L., J.L., K.O., D.C., L.F., M.A., J.A.S., E.E.F., I.L., R.D., B.P., J.P.M., D.A.R., J.A.C., S.O., A.Z., C.E.H., R.B., J.A.-S., J.M. and I.L.Y. collected field data.

Competing interests. We declare we have no competing interests.

Funding. We acknowledge and thank the Southwest Fisheries Science Center (NMFS-NOAA) for processing and analytical support, as well as the Galapagos Science Center/Universidad San Francisco de Quito for supplementary laboratory processing of samples from Ecuador. We recognize the financial contributions from several organizations that enabled the operation of the hawksbill conservation projects during which the samples presented in this paper were collected, including the National Fish and Wildlife Foundation (NFWF), US Fish & Wildlife Service (USFWS), International Seafood Sustainability Foundation (ISSF), William H. Donnor Foundation and SeeTurtles.org. This research was also funded in part by a grant from the Pacific Rim Research Program of the University of California.

Acknowledgements. We thank the following individuals: Brad Nahill, Hoyt Peckham, Neftali Sanchez, Obed Rivera, Luis Manzanera, Liza Gonzalez, Jacinto Rodriguez, Luis Mera, Ricardo González, Gabriela Serra-Valente, Amy Lanci, Erin LaCasella and Amy Frey. We recognize the following groups and organizations for logistical support: Hoteles de Guayabitos, ?!Careyes Foundation, Comité Carey de la Bahía de Jiquilisco, Comité Carey de la Reserva Natural Estero Padre Ramos, Fundarrecife, Manta Raya Hospedaje, Machalilla National Park and The Ocean Foundation for additional logistical support.

References

- Ricklefs RE. 1987 Community diversity: relative roles of local and regional processes. *Science* **235**, 167–171. (doi:10.1126/science.235.4785.167)
- Tilman D, Kareiva P. 1997 *Spatial ecology: the role of space in population dynamics and interspecific interactions*, p. 416. Princeton, NJ: Princeton University Press.
- Pritchard PCH. 1976 Post-nesting movements of marine turtles (*Cheloniidae* and *Dermochelyidae*) tagged in the Guianas. *Copeia* **4**, 749–754. (doi:10.2307/1443458)
- Godley BJ, Blumenthal JM, Broderick AC, Coyne MS, Godfrey MH, Hawkes LA, Witt MJ. 2008 Satellite tracking of sea turtles: where have we been and where do we go next? *Endang. Species Res.* **4**, 3–22. (doi:10.3354/esr00060)
- Carr A. 1987 New perspectives on the pelagic stage of sea turtle development. *Conserv. Biol.* **1**, 103–121. (doi:10.1111/j.1523-1739.1987.tb00020.x)
- Bolten AB. 2003 Active swimmers—passive drifters: the oceanic juvenile stage of loggerheads in the Atlantic system. In *Loggerhead head sea turtles* (eds AB Bolten, BE Witherington), pp. 63–78. Washington, DC: Smithsonian Books.
- Luschi P, Hays GC, Papi F. 2003 A review of long-distance movements by marine turtles, and the possible role of ocean currents. *OIKOS* **103**, 293–302. (doi:10.1034/j.1600-0706.2003.12123.x)
- Lohmann KJ, Putman NF, Lohmann CMF. 2012 The magnetic map of hatching loggerhead sea turtles. *Curr. Opin. Neurobiol.* **22**, 336–342. (doi:10.1016/j.conb.2011.11.005)
- Bolten AB. 2003 Variation in sea turtle life history patterns: neritic vs. oceanic developmental stages. In *The biology of sea turtles*, vol. II (eds PL Lutz, J Musick, J Wyneken), pp. 243–257. Boca Raton, FL: CRC Press.
- Avens L, Snover M. 2013 Age and age estimation in sea turtles. In *The biology of sea turtles*, vol. III (eds J Wyneken, KJ Lohmann, JA Musick), pp. 97–133. Boca Raton, FL: CRC Press.
- Jensen MP, FitzSimmons NN, Dutton PH. 2013 Molecular genetics of sea turtles. In *The biology of sea turtles*, vol. III (eds J Wyneken, KJ Lohmann, JA Musick), pp. 135–162. Boca Raton, FL: CRC Press.
- Blumenthal JM *et al.* 2009 Turtle groups or turtle soup: dispersal patterns of hawksbill turtles in the Caribbean. *Mar. Ecol.* **18**, 4842–4853. (doi:10.1111/j.1365-294X.2009.04403.x)
- Hays GC, Fossette S, Katselidis KA, Mariani P, Schofield G. 2010 Ontogenetic development of migration: Lagrangian drift trajectories suggest a new paradigm for sea turtles. *J. R. Soc. Interface* **7**, 1319–1327. (doi:10.1098/rsif.2010.0009)
- Gaspar P, Benson SR, Dutton PH, Réveillére A, Jacob G, Meetoo C, Dehecq A, Fossette S. 2012 Oceanic dispersal of juvenile leatherback turtles: going beyond passive drift modeling. *Mar. Ecol. Prog. Ser.* **457**, 265–284. (doi:10.3354/meps09689)
- Putman NF, Mansfield KL. 2015 Direct evidence of swimming demonstrates active dispersal in the sea

- turtle 'lost years'. *Curr. Biol.* **4**, 1221–1227. (doi:10.1016/j.cub.2015.03.014)
16. Bowen BW, Bass AL, Soares LS, Toonen RJ. 2004 Natal homing in juvenile loggerhead turtles (*Caretta caretta*). *Mol. Ecol.* **13**, 3797–3808. (doi:10.1111/j.1365-294X.2004.02356.x)
 17. Bowen BW, Grant WS, Hillis-Starr Z, Shaver DJ, Bjorndal KA, Bolten AB, Bass AL. 2007 Mixed stock analysis reveals the migrations of juvenile hawksbill turtles (*Eretmochelys imbricata*) in the Caribbean Sea. *Mol. Ecol.* **16**, 49–60. (doi:10.1111/j.1365-294X.2006.03096.x)
 18. Bass AL, Epperly SP, Braun-McNeill J. 2004 Multi-year analysis of stock composition of a loggerhead turtle (*Caretta caretta*) foraging habitat using maximum likelihood and Bayesian methods. *Conserv. Genet.* **5**, 783–796. (doi:10.1007/s10592-004-1979-1)
 19. Naro-Maciel E, Becker JH, Lima EHS, Marcovaldi MA and Desalle R. 2007 Testing dispersal hypotheses in foraging green sea turtles (*Chelonia mydas*) of Brazil. *J. Hered.* **98**, 29–39. (doi:10.1093/jhered/esl050)
 20. Naro-Maciel E, Bondioli ACV, Martin M, Almeida AP, Baptistotte C, Bellini C, Marcovaldi MA, Santos AJB, Amato G. 2012 The interplay of homing and dispersal in green turtles: a focus on the southwestern Atlantic. *J. Hered.* **103**, 792–805. (doi:10.1093/jhered/ess068)
 21. Monzón-Arguello C, López-Jurado LF, Rico C, Marco A, López P, Hays GC, Lee PLM. 2010 Evidence from genetic and Lagrangian drifter data for transatlantic transport of small juvenile green turtles. *J. Biogeogr.* **37**, 1752–1766. (doi:10.1111/j.1365-2699.2010.02326.x)
 22. Prosdocimi L, Carman VG, Albareda DA, Remis MI. 2011 Genetic composition of green turtle feeding grounds of coastal waters of Argentina based on mitochondrial DNA. *J. Exp. Mar. Biol. Ecol.* **412**, 37–45. (doi:10.1016/j.jembe.2011.10.015)
 23. Bjorndal KA. 1982 *Biology and conservation of sea turtles*. Washington, DC: Smithsonian Institution Press.
 24. Gaos AR *et al.* 2010 Signs of hope in the eastern Pacific: international collaboration reveals encouraging status for a severely depleted population of hawksbill turtles *Eretmochelys imbricata*. *Oryx* **44**, 595–601. (doi:10.1017/S0030605310000773)
 25. Wallace BP *et al.* 2011 Regional management units for marine turtles: a novel framework for prioritizing conservation and research across multiple scales. *PLoS ONE* **5**, e15465. (doi:10.1371/journal.pone.0015465)
 26. Gaos AR *et al.* 2012 Spatial ecology of critically endangered hawksbill turtles *Eretmochelys imbricata*: implications for management and conservation. *Mar. Ecol. Prog. Ser.* **450**, 181–194. (doi:10.3354/meps09591)
 27. Gaos AR, Lewison RL, Wallace B, Yañez IL, Liles M, Baquero A, Seminoff JA. 2012 Dive behaviour of adult hawksbills (*Eretmochelys imbricata*, Linnaeus 1766) in the eastern Pacific Ocean highlights shallow depth-use by the species. *J. Exp. Mar. Biol. Ecol.* **432–433**, 171–178. (doi:10.1016/j.jembe.2012.07.006)
 28. Carrión-Cortez J, Canales-Cerro C, Arauz R, Riosmena-Rodríguez R. 2013 Habitat use and diet of juvenile eastern Pacific hawksbill turtles (*Eretmochelys imbricata*) in the north Pacific coast of Costa Rica. *Chelonian Conserv. Biol.* **12**, 235–245. (doi:10.2744/CCB-1024.1)
 29. Gaos AR *et al.* 2016 Hawksbill turtle terra incognita: conservation genetics of eastern Pacific rookeries. *Ecol. Evol.* **6**, 1251–1264. (doi:10.1002/ece3.1897)
 30. Excoffier L, Lischer HE. 2010 Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* **10**, 564–567. (doi:10.1111/j.1755-0998.2010.02847.x)
 31. Peakall R, Smouse PE. 2006 genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol. Ecol. Notes* **6**, 288–295. (doi:10.1111/j.1471-8286.2005.01155.x)
 32. Pella J, Masuda M. 2001 Bayesian methods for analysis of stock mixtures from genetic characters. *Fish. Bull.* **99**, 151–167.
 33. Gaos A *et al.* 2017 Living on the edge: hawksbill turtle nesting and conservation along the eastern Pacific rim. *Lat. Am. J. Aquat. Res.* **45**, 572–584. (doi:10.3856/vol45-issue3-fulltext-7)
 34. Thums M, Whiting SC, Reisser JW, Pendoley KL, Pattiaratchi CB, Harcourt RG, McMahon CR, Meekan MG. 2013 Tracking sea turtle hatchlings—a pilot study using acoustic telemetry. *J. Exp. Mar. Biol. Ecol.* **440**, 156–163. (doi:10.1016/j.jembe.2012.12.006)
 35. Bowen BW, Bass AL. 1997 Movement of hawksbill turtles: what scale is relevant to conservation, and what scale is resolvable with mtDNA data? *Chelonian Conserv. Biol.* **2**, 440–442.
 36. Meylan AB, Bowen BW, Avise JC. 1990 A genetic test of the natal homing versus social facilitation models for green turtle migration. *Science* **248**, 724–727. (doi:10.1126/science.2333522)
 37. Mansfield KL, Wyneken J, Porter WP, Luo J. 2014 First satellite tracks of neonate sea turtles redefine the 'lost years' oceanic niche. *Proc. R. Soc. B* **281**, 20133039. (doi:10.1098/rspb.2013.3039)
 38. Lohmann KJ, Lohmann CMF, Brothers JR, Putman NF. 2013 Natal homing and imprinting in sea turtles. In *The biology of sea turtles*, vol. III (eds J Wyneken, KJ Lohmann, JA Musick), pp. 59–78. Boca Raton, FL: CRC Press.
 39. Davis JM, Stamps JA. 2004 The effect of natal experience on habitat preference. *Trends Ecol. Evol.* **19**, 411–416. (doi:10.1016/j.tree.2004.04.006)
 40. Brothers JR, Lohmann KJ. 2015 Evidence for geomagnetic imprinting and magnetic navigation in the natal homing of sea turtles. *Curr. Biol.* **25**, 392–396. (doi:10.1016/j.cub.2014.12.035)
 41. Liles MJ *et al.* 2017 Survival on the rocks: high bycatch in lobster gillnet fisheries threatens hawksbill turtles on rocky reefs along the Eastern Pacific coast of Central America. *Lat. Am. J. Aquat. Res.* **45**, 521–539. (doi:10.3856/vol45-issue3-fulltext-3)
 42. Kessler WS. 2006 The circulation of the eastern tropical Pacific: a review. *Prog. Oceanogr.* **69**, 181–217. (doi:10.1016/j.pocean.2006.03.009)
 43. Gaos AR *et al.* 2012 Shifting the life-history paradigm: discovery of novel habitat use by hawksbill turtles. *Biol. Lett.* **8**, 54–56. (doi:10.1098/rsbl.2011.0603)
 44. Liles MJ *et al.* 2015 One size does not fit all: importance of adjusting conservation practices for endangered hawksbill turtles to address local nesting habitat needs in the eastern Pacific Ocean. *Biol. Conserv.* **184**, 405–413. (doi:10.1016/j.biocon.2015.02.017)
 45. Chung FC, Pilcher NJ, Salmon M, Wyneken J. 2009 Offshore migratory activity of hawksbill turtle (*Eretmochelys imbricata*) hatchlings, II. Swimming gaits, swimming speed, and morphological comparisons. *Chelonian Conserv. Biol.* **8**, 35–42. (doi:10.2744/CCB-0716.1)
 46. Wildermann N, Critchell K, Fuentes MMPB, Limpus CJ, Wolanski E, Hamann M. 2017 Does behaviour affect the dispersal of flatback post-hatchlings in the Great Barrier Reef? *R. Soc. open sci.* **4**, 170164. (doi:10.1098/rsos.170164)
 47. Walker TA, Parmenter CJ. 1990 Absence of a pelagic phase in the life cycle of the flatback turtle, *Natator depressa* Garman. *J. Biogeogr.* **17**, 275–278. (doi:10.2307/2845123)
 48. Clutton-Brock TH. 1988 *Reproductive success: studies of individual variation in contrasting breeding systems*, p. 538. Chicago, IL: University of Chicago Press.
 49. Shillinger GL *et al.* 2008 Persistent leatherback turtle migrations present opportunities for conservation. *PLoS Biol.* **6**, e171. (doi:10.1371/journal.pbio.0060171)
 50. Mortimer JA, Meylan PA, Donnelly M. 2007 Whose turtles are they, anyway? *Mol. Ecol.* **16**, 17–18. (doi:10.1111/j.1365-294X.2006.03252.x)
 51. Whiting SD, Murray W, Macrae I, Thorn R, Chongkin M, Koch AU. 2008 Non-migratory breeding by isolated green sea turtles (*Chelonia mydas*) in the Indian Ocean: biological and conservation implications. *Naturwissenschaften* **95**, 355–360. (doi:10.1007/s00114-007-0327-y)
 52. McClenahan L, Jackson JBC, Newman MJH. 2006 Conservation implications of historic sea turtle nesting beach loss. *Front. Ecol. Environ.* **4**, 290–296. (doi:10.1890/1540-9295(2006)4[290:CIOHST]2.0.CO;2)
 53. Warner RP, Chesson PL. 1985 Coexistence mediated by recruitment fluctuations: a field guide to the storage effect. *Am. Nat.* **125**, 769–787. (doi:10.1086/284379)
 54. Hendry AP, Castric V, Kinnison MT, Quinn TP. 2004 *The evolution of philopatry and dispersal: homing versus straying in salmonids*. In *Evolution illuminated: salmon and their relatives* (eds AP Hendry, SC Stearns), pp. 52–90. New York, NY: Oxford University Press.
 55. Coulson J. 2016 A review of philopatry in seabirds and comparisons with other waterbird species. *Waterbirds* **39**, 229–240. (doi:10.1675/063.039.0302)
 56. Gaos AR *et al.* 2017 Data from: Natal foraging philopatry in eastern Pacific hawksbill turtles. Dryad Digital Repository. (<http://dx.doi.org/10.5061/dryad.38123>)