FILE COPY

# NOAA Technical Memorandum NMFS



**DECEMBER 1982** 

# REVISED UPDATE AND RETRIEVAL SYSTEM FOR THE CALCOFI OCEANOGRAPHIC DATA FILE

Lawrence E. Eber and Nancy Wiley



NOAA-TM-NMFS-SWFC-24

U.S. DEPARTMENT OF COMMERCE National Oceanic and Atmospheric Administration National Marine Fisheries Service Southwest Fisheries Center

#### **NOAA Technical Memorandum NMFS**

The National Oceanic and Atmospheric Administration (NOAA), organized in 1970, has evolved into an agency which establishes national policies an manages and conserves our oceanic, coastal, and atmospheric resources. An organizational element within NOAA, the Office of Fisheries is responsible for fisheries policy and the direction of the National Marine Fisheries Service (NMFS).

In addition to its formal publications, the NMFS uses the NOAA Technical Memorandum series to issue informal scientific and technical publications when complete formal review and editorial processing are not appropriate or feasible. Documents within this series, however, reflect sound professional work and may be referenced in the formal scientific and technical literature.

# **NOAA Technical Memorandum NMFS**



This TM series is used for documentation and timely communication of preliminary results, interim reports, or special purpose information; and have not received complete formal review, editorial control, or detailed editing.

**DECEMBER 1982** 

# REVISED UPDATE AND RETRIEVAL SYSTEM FOR THE CALCOFI OCEANOGRAPHIC DATA FILE

Lawrence E. Eber Southwest Fisheries Center National Marine Fisheries Service La Jolla, CA 92038

and

Nancy Wiley
Southwest Fisheries Center
National Marine Fisheries Service
Tiburon, CA 94920

NOAA-TM-NMFS-SWFC-24

U.S. DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary
National Oceanic and Atmospheric Administration
John V. Byrne, Administrator
National Marine Fisheries Service
William G. Gordon, Assistant Administrator for Fisheries

# REVISED UPDATE AND RETRIEVAL SYSTEM FOR THE CALCOFI OCEANOGRAPHIC DATA FILE

#### L.E. Eber and Nancy Wiley

#### **PREFACE**

This report supersedes Administrative Report No. LJ-76-16 of the Southwest Fisheries Center (SWFC), entitled "A Retrieval and Display System for the CalCOFI Oceanographic Data File," by Eber and Wiley. The need for revision of that report arose from the fact that most of the retrieval options described therein are no longer available owing to substantive modifications in the retrieval software.

Certain portions of the earlier report containing background information about the CalCOFI program and procedures used in assembling the SWFC version of the oceanographic data file have been carried over to this revision.

#### INTRODUCTION

Oceanographic observations have been made over an extensive grid of stations in the California Current region since 1949. The grid was designed during development of a systematic sampling program to determine the major spawning areas of the Pacific sardine off the coasts of the United States and Baja California, Mexico. With the demise of the sardine, emphasis in the program was redirected toward the Northern Anchovy.

Most of the work involved in collection, processing and analysis of these observations was performed by Scripps Institution of Oceangraphy (SIO), National Marine Fisheries Service (NMFS) and California Department of Fish and Game (CF&G), (Kramer, et al. 1972).

The physical and chemical data from CalCOFI surveys have been compiled by SIO into data reports (listed in Appendix 1), which include values of temperature, salinity and oxygen tabulated at observed and at standard depths, and computed values of sigma-t, thermosteric anomaly and geopotential anomaly at standard depths. Some observations of phosphorus, nitrite, nitrate and silicon are also included. Physical and chemical data for the years 1949 through 1959 were also published in Oceanic Observations of the Pacific.1

These data have been used by many investigators working on problems associated with the California Current. Some of the data have been presented graphically as horizontal and vertical sections in the CalCOFI Atlas Series

 $<sup>^{1}</sup>$ Oceanic Observations of the Pacific, published by SIO, is a series of annual volumes of oceanographic data collected in the Pacific Ocean and its adjacent seas.

(Anom., 1963; Wyllie, J.G., 1966; Wyllie, J.G. and R.J. Lynn, 1971; Thomas, W.H. and D.L.R. Seibert, 1974; Eber, L.E., 1977; Lynn, R.J., K.A. Bliss and L.E. Eber, 1982). There remains, however, a large amount of untapped information in the CalCOFI data resource which warrants the development of a system for convenient and economical retrieval.

#### REASSEMBLY OF THE CALCOFI OCEANOGRAPHIC FILE

Oceanographic data were processed initially by the Data Collection and Processing Group (DCPG) at SIO and subsequently archived at the National Oceanographic Data Center (NODC). The NODC format for oceanographic station data is a BCD card-image format extended to 120 characters. An oceanographic station is represented by a master "card" containing station identification and surface observations followed by detail "cards" containing subsurface data at observed or standard (interpolated) depths.

The Automatic Data Processing unit at SWFC obtained, from SIO, copies of CalCOFI oceanographic data on magnetic tape in the NODC format, for the period 1950-1968. The data were partitioned into 5 sections of the CalCOFI station pattern and ordered by year, month and station number within each section.

The first step taken in reassembling this file was to transfer the data to the Burroughs B6700 disk at which time the 120 character format was shortened to 90 characters, by deleting certain fields not required by SWFC. Then the file was subjected to an editing procedure in which format, sequence, syntax and identification errors were diagnosed and corrected. This process did not include an inspection of the parameter values (temperature, salinity, etc.). When the editing was completed the 5 sections were merged and the file was resorted into chronological order by cruise code.

Data retrieval from large disk or tape files in character (card-image) formats can be slow and inefficient. Consequently, a packed binary format was designed which would reduce the size of the oceanographic file by about 75%. A description of this format is given in Appendix 2. An ALGOL program, PACKOCEAN, was written and used to convert the abridged NODC format to the new packed binary format and to create a new oceanographic file containing all the cruises from 1950 through 1968.

In the next decade, issuance of fully processed oceanographic data from ongoing CalCOFI cruises began to lag. To assuage the needs of SWFC for current data, an arrangement was made with DCPG to obtain copies of the oceanographic data from completed cruises at a preliminary stage of the processing. The data were obtained in a card format used by DCPG and transferred to B6700 disk files. These data did not include the computed density parameters (sigma-t, etc.) or interpolations to standard depths. Consequently, a copy of the FORTRAN program used by DCPG to perform these computations was procured, translated into ALGOL and incorporated, along with much of PACKOCEAN, into a program named OCEANDATA/UPDATE. This program has been used to convert the DCPG card format to the SWFC packed binary format and update the CalCOFI oceanographic file with data from cruises in the period 1969 through 1978.

#### THE CALCOFI STATION PLAN - DATA COVERAGE

The CalCOFI station pattern is shown in Figure 1. It was based on line 80, which extends seaward off Point Conception, California in a direction 30 degrees south of west. This is approximately perpendicular to the California coastline between Point Conception and Cape Mendocino. Other principal, or cardinal, lines were laid out parallel to line 80 on either side at intervals of 120 nautical miles. The added lines were numbered using increments of 10, increasing southward and decreasing northward.

Two intermediate, or ordinal, lines were inserted between each pair of cardinal lines, at intervals of 40 nautical miles. This corresponds to a line number increment of 3.333..., which results in fractional line numbers for all ordinal lines. By convention, however, ordinal line numbers have been rounded off to end in 3's and 7's.

Stations on CalCOFI lines were established with reference to a perpendicular through line 80, intersecting at a point about 40 nautical miles off the coast. This intersection of the perpendicular with line 80 was designated station 60, as were its intersections with each of the other lines, both cardinal and ordinal. Other stations on each line were laid out initially at intervals of 40 nautical miles, numbered by increments of 10, increasing in the offshore direction. This scheme provided for insertion of intermediate stations at intervals as small as 4 nautical miles without requiring fractional increments between station numbers.

Conventional station codes were made up of a line number and a station number, separated by a decimal point. For example, 80.60 refers to station 60 on line 80. For computer processing the conventional code was modified to a six digit integer (with leading zero optional) composed of a 3 digit line number followed by a 3 digit station number. Thus, station code 80.60 would be changed to either 080060 or 80060. Station codes in the SWFC version of the CalCOFI oceanographic data file are in the modified form.

The station pattern shown in Figure 1 has been used in planning CalCOFI surveys since 1950. Coverage of the pattern by survey cruises has varied considerably over the years owing to changes in sampling strategy, availability of ship time and other factors. In general, the cruises were conducted monthly during the 1950's and quarterly during the 1960's. Since 1969, cruises have been limited to every third year.

Table 1 contains the station codes for all CalCOFI stations which were occupied on six or more cruises in the period 1950-1978. The geographical location is given for each station, along with a list of the cruises on which it was occupied.

Recent innovations in sampling strategy devised particularly for anchovy eggs involve higher spacial resolution, and require greater precision in designating station locations than was possible under the convention of rounding off line and station numbers to the nearest integer. Consequently, CalCOFI line and station numbers are presently specified to one decimal place

for purposes of laying out cruise tracks. A computer procedure derived by Eber and Hewitt (1979) has been implemented to facilitate determination of latitude and longitude of any station specified with fractional values for line and station number.

# DATA RETRIEVAL FROM THE OCEANOGRAPHIC DATA FILE - CCFIHYDRO

Software for retrieval of CalCOFI data from the SWFC packed binary file provides a variety of retrieval options, referred to here as data selection modes, or run modes. Designation of these options was based both on anticipated application requirements and on the data format in the packed binary file. Each record in this file represents one CalCOFI station and consists of three parts: (1) surface data which includes time and location identification fields and wind, weather and sea state observations, (2) physical and chemical parameters at observed depths, and (3) interpolated physical and chemical parameters and computed density parameters at standard depths.

An ALGOL program, CCFIHYDRO, was designed to extract data from the packed binary file in accordance with selection criteria based on station codes, cruise codes and standard or observed depths. The output produced by this program can be in one of three forms: (1) a subfile of selected records copied intact from the packed binary source file, (2) a working file containing observed and/or interpolated values of subsurface physical and chemical parameters at selected stations and/or cruises, (3) a printout of the surface data (surface observations and identification fields) for selected stations and/or cruises.

Data selection in CCFIHYDRO is controlled by one or more sets of data selection cards containing station lists, cruise lists or depth lists which are submitted at run time. Selection modes which create subfiles are the most efficient in terms of computer time but allow only one kind of selection criteria, either stations or cruises. A subfile created by one run of CCFIHYDRO can be used as input for subsequent runs.

Working files are created under selection modes for which all three criteria--stations, cruises and depths--may be required. Data in these files are organized by cruises and contain values for six parameters: temperature, salinity, oxygen, sigma-t, thermosteric anomaly and geopotential anomaly along with CalCOFI station number, year, month, day and depth. These files are written in a simple binary format which can be readily converted to a cardimage (EBCDIC) or ASCII format for transfer to tape or to another computer.

Data selection modes which involve surface data produce only printer output and are provided as a means of indexing selected segments of the oceanographic file and displaying weather and sea conditions at times when subsurface observations were taken.

CCFIHYDRO incorporates an external procedure, UNPACKV, for actual extraction and unpacking of the bit-fields which store parameter values in the packed binary file. UNPACKV is efficient, involving no conversions from one

number base to another, and provides for extraction of any parameter in the source file. It can be linked to any ALGOL or FORTRAN main program to perform retrieval functions outside the scope of CCIFHYDRO. Documentation on CCFIHYDRO and UNPACKV is given in Appendices 3 and 4.

Three ancillary programs written for manipulation of workfiles created by CCFIHYDRO are MAKOCNSUB, ADDSTN and PRTOCNDAT. MAKOCNSUB extracts data from workfiles using either cruises, stations or depths as selection criteria. The extracted data are put into smaller workfiles. This program provides an alternate way of selecting data for workfiles which might be used as input for contouring by EDMAP2. ADDSTN merges two workfiles, producing a third workfile in which the merged station codes are stored in ascending order within cruises just as in the two input files. PRTOCNDAT copies workfiles to the printer, or to disk or tape. In each case the simple binary format is converted to a character format (EBCDIC OR ASCII). One or more of these output options may be specified at run time. Documentation on MAKOCNSUB, ADDSTN, PRTOCNDAT and six additional utility programs is given in Appendix 5.

The specifications and options which set the data selection modes in CCFIHYDRO are submitted at run time by means of a run specification card and one or more sets of data selection cards. In the next two sections these control cards are described in detail.

#### CCFIHYDRO RUN SPECIFICATION CARD

The run specification card has six fields. The first field is reserved for a user title, which must be entered in cols. 1-24. The title may be any EBCDIC (or ASCII) string. The first 12 cols. of the title will be copied into identification records of workfiles created during the run. This provides a means of identifying outputs.

The remaining five fields are entered in free-field format, using one or more spaces as field delimiters. The symbol (#) terminates processing of the run specification card and may be used at any point after the fourth field. Default specifications will be assigned for the remaining fields.

Valid entries for the run specification card are listed in Table 2. Any entry in fields 2, 3 or 4 which does not conform to this table will result in program termination. Unrecognized entries in any other field will result in assignment of the default value for that field. The specifications in fields 2, 3 and 4 determine the data selection mode for a particular run. Valid combinations of entries in these fields are shown in Table 3. Each mode imposes restrictions on the choice of entries for field 5 and on the contents of data selection cards. The program does not check whether specifications in fields 2-4 constitute a valid combination. Consequently, one which does not conform to Table 3, may produce unpredictable results.

<sup>2</sup>EDMAP2 is an ALGOL program for analysis and contouring of environmental data fields on a two dimensional grid. Documentation on this program is contained in NOAA Tech. Memo. NMFS-SWFC-18, April 1982.

Data selection modes comprise three groups, each associated with a different kind of output. In the first six modes (Table 3) the only elements extracted from the source file are the station data, including ship's position, station code, date, time and surface weather and sea state observations. This information is output to a printer in tabular format. In the next nine modes (7-15) six subsurface data parameters are extracted and output to a disk file in a simple binary format. In the last two modes (17-18) complete records are extracted from the source file and copied to a subfile on disk without change of format, and key identification fields from the records are printed.

The second group of data selection modes (7-15) were intended to provide flexible selection criteria for retrieval from the source file and to facilitate mapping the oceanographic parameters on horizontal and vertical sections, using the analysis and contouring program EDMAP2. For this purpose, the specification in field 2 is interpreted as the Y-coordinate and that in field 3, as the X-coordinate of a two-dimensional grid representing the section. For the purpose of simply extracting data from the source file to create a working file, however, many of the data selection modes are redundant, and most data selection requirements can be satisfied with modes 7 or 8 for standard depths and modes 13 or 14 for observed depths. Descriptions of valid specification parameters for each field are given below.

# Field 1 (Cols. 1-24):

(Title) Any character string. Cols. 1-12 will be copied into identification records of output files in run modes 7-15.

#### Field 2:

DEPTH Specifies standard or observed depths as 1st criteria (1st set of data selection cards) for data retrieval in run modes 7-15. Also designates depth as the Y-coordinate for mapping.

DISTN Specifies station codes as the 1st criteria for data retrieval in run modes 7-15. Also designates distance normal to the coast (along CalCOFI lines) as the Y-coordinate for mapping.

DISTP Specifies station codes as the 1st criteria for data retrieval in run modes 7-15. Also designates distance parallel to the coast (across CalCOFI lines) as the Y-coordinate for mapping.

SUBFIL Specifies that a subfile is to be created by copying records intact from the source (packed binary) file, using either station codes or cruise codes as selection criteria as specified in Field 3.

#### Field 3

TIME Specifies cruise codes as the 2nd criteria (2nd set of data selection cards) for data retrieval in run modes 7-15, and the sole

criteria in run mode 18. Also designates time as the X-coordinate for mapping.

DISTN Specifies station codes as the 2nd criteria for data selection in run modes 1-15. Also designates distance normal to the coast (along CalCOFI lines) as the X-coordinate for mapping.

DISTP Specifies station codes as the 2nd criteria for data selection in run modes 1-15. Also designates distance parallel to the coast (across CalCOFI lines) as the X-coordinate for mapping.

STATIONS Specifies station codes as the sole criteria for data selection in run mode 17. It may be used in place of either DISTN or DISTP for other modes when the alignment of the stations to be selected has no special significance.

CRUISES Specifies cruise codes as the sole criteria for data selection in run mode 18. It may be used in place of TIME under all circumstances.

#### Field 4:

STAT Specifies that surface data will be extracted from selected source file records and printed. The surface elements include ship's position, date and time of observation, depth to bottom, wind, weather and sea state. Run mode will be set in the range 1-6 depending on entries in Fields 2 and 3.

STD Specifies that temperature, salinity, oxygen, sigma-t, thermosteric anomaly and geopotential anomaly at standard depths will be extracted from selected source records and stored in a working file (OCNDAT) on disk. Run mode will be set in the range 7-12, depending on entries in Fields 2 and 3.

OBS, OBSTD Specifies that temperature, salinity, oxygen and sigma-t at observed depths, or at observed and standard depths, will be extracted from selected source file records and stored in a working file (OCNDAT) on disk. Run mode will be set in the range 13-15, depending on entries in Fields 2 and 3.

ALL Specifies that selected source records will be copied intact into a subfile (SUBFIL) on disk. Run mode will be set to 17 or 18 depending on the entry in Field 3. The entry in Field 2 must be SUBFIL.

#### Field 5:

(Cruise code) Designates a single cruise for data selection from the source file.

(Station code) Designates a single station for data selection from the source file.

(Std Depth) Designates a single standard depth for data selection from the source file.

O Specifies either that all cruises are to be accepted for a station

list submitted as the 2nd set of data selection cards (all odd-numbered modes), or that all stations are to be accepted for a cruise list submitted as the 2nd set of data selection cards (run modes 2, 8 or 14), or that data will be extracted from selected source records only for standard depth = 0 meters (run modes 10 and 12).

1 Specifies that a cruise list is being submitted as the 3rd set of data selection cards (odd-numbered modes only).

#### Field 6:

CONCAT Causes records selected from the source file to be concatenated with an existing subfile on disk. Run modes 17 and 18.

PRTREC Causes all fields in records selected from the source file to be printed in a format similar to that used in the SIO Physical and Chemical Data Reports (Appendix 1).

WRTREC Causes all fields in records selected from the source file to be written to a file (tape or disk) in ASCII code.

PRWREC Causes the actions described for both PRTREC and WRTREC to occur.

# Terminates processing of the run specification card. It can be entered anytime after Field 4: default values will be assigned to unspecified Fields. The default for Field 5 is 0 and for Field 6 is NOT CONCAT.

#### CCFIHYDRO DATA SELECTION CARDS

One or more sets of data selection cards must be submitted with each run. They are to be inserted just after the run specification card in the Job set-up (Appendix 3). Data selection cards contain lists of depths, station codes, cruise codes or consecutive sequence numbers. The kind(s) of data selection cards appropriate for each run mode are indicated in Table 3.

Data selection cards must be submitted in a fixed-field format given by

(10(X1, I6, X1)) for ALGOL syntax.

The first field (cols. 2-7) on the first card of each set must contain the number of subsequent entries submitted in that set. For example, if a set of data selection cards contains 27 station codes then the number in the first field of the first card in the set would be 27.

For run modes 1, 2, 3, 5 and 11 the 1st set of data selection cards must have a single entry, which will be either 0 (run modes 1, 2, 3, 5) or a standard depth value (run mode 11).

Stations should be entered on data selection cards as 5 or 6 digit station codes in either ascending or descending order. Cruises should be entered as 4 digit year-month codes in chronological order. Standard depths

should be entered in order of increasing depth. Sequence numbers (run modes 13-15) should be entered as consecutive integers starting with 1. They are used by CCFIHYDRO as array indices when storing observed depth values extracted from source records. The highest sequence number submitted determines the maximum number of observed depths for which data will be extracted.

A maximum of 300 entries is allowed for each set of data selection cards.

#### CCFIHYDRO SOURCE FILES AND OUTPUT FILES

CCFIHYDRO uses four files in the performance of its data retrieval function. The internal names are CCFI, SUBFIL, OCNDAT and CALTAP. These are equated to directory names of physical files on disk or tape at run time.

CCFI is the CalCOFI source file in the SWFC packed binary format. It must always be available for input when CCFIHYDRO is run.

SUBFIL is one of the output file option created in run modes 17-18. It has the same format as CCFI. A SUBFIL created in one run can be re-designated as a source file (CCFI) for subsequent runs.

CALTAP is a second output file options created in run modes 17-18. It will contain all of the information available in the source file (CCFI) for specified stations or cruises, in ASCII format. This option is invoked when either WRTREC or PRWREC are entered in the sixth field of the run specification card.

OCNDAT is the output file created in run modes 7-15. It will contain one or more data sets, each consisting of a header, or identification, record, followed by a varying number of data records. OCNDAT files created under odd-numbered run modes will have one data set for each selected cruise. Under even-numbered modes there will be one data set for each of one or more sequences of cruises as specified by data selection cards.

The data records in a data set each contain nine fields. The first field will have either station codes (for odd-numbered run modes) or cruise codes (for even-numbered run modes). The second field will contain a 6-digit code representing a year-month-day code (YYMMDD). The third field will contain depth in meters. The remaining six fields will contain temperature, salinity, oxygen, sigma-t, thermosteric anomaly and geopotential anomaly, respectively. A zero value for any parameter (except geopotential anomaly at 500 meters) denotes missing data.

OCNDAT files can be submitted as input to the analysis and contouring program EDMAP2 for making vertical sections (depth vs. distance or time) or horizontal sections at standard depths, for any of the six data fields.

#### REFERENCES

- Anonymous. 1963. CalCOFI Atlas of 10-meter temperatures and salinities 1949 through 1959. Calif. Coop. Oceanic Fish. Invest., Atlas. No. 1, DCPG, MLR Prog., SIO.
- Eber, L.E. 1977. Contoured depth-time charts (0 to 200 m, 1950 to 1966) of temperature, salinity, oxygen and sigma-t at 23 CalCOFI stations in the California Current. Calif. Coop. Oceanic Fish. Invest., Atlas No. 25, DCPG, MLR Prog., SIO.
- Kramer, D., M.J. Kalin, E.G. Stevens, J.R. Thrailkill and J.R. Zweifel. 1972. Collecting and processing data in fish eggs and larvae in the California Current Region. NOAA Tech. Rept. NMFS Circ. 370, 38 p.
- Lynn, R.J., K.A. Bliss and L.E. Eber. 1982. Vertical and horizontal distributions of seasonal mean temperature, salinity, sigma-t, stability, dynamic height, oxygen and oxygen saturation in the California Current, 1950-1978. Calif. Coop. Oceanic Fish. Invest., Atlas No. 30, MLR Prog., SIO.
- Thomas, W.H. and D.L.R. Seibert. 1974. Distribution of nitrate, nitrite, phosphate and silicate in the California Current region, 1969. Calif. Coop. Oceanic Fish. Invest., Atlas No. 20, DCPG, MLR Prog., SIO.
- Wyllie, J.G. 1966. Geostrophic flow of the California Current at the surface and at 200 meters. Calif. Coop. Oceanic Fish. Invest., Atlas No. 4, DCPG, MLR Prog., SIO.
- Wyllie, J.G. and R.J. Lynn. 1971. Distribution of temperature and salinity at 10 meters, 1960-69 and mean temperature, salinity and oxygen at 150 meters, 1950-1968 in the California Current. Calif. Coop. Oceanic Fish. Invest., Atlas No. 15, DCPG, MLR Prog., SIO.

Appendix 1. List of Physical and Chemical Data Reports from MLR and CalCOFI-Cruises, Issued By Scripps Institution of Oceanography.

Research cruises in the California Current region conducted by the Marine Life Research (MLR) Program of SIO, in cooperation with the U.S. Fish and Wildlife Service, were initiated in March 1949. The original survey pattern and station numbering scheme was abandoned after the first 10 MLR cruises, at which time the pattern shown in Figure 1 was adopted. The Physical and Chemical Data Reports continued to bear MLR cruise numbers until May 1952, when the CalCOFI cruise designations were added. As indicated in the Introduction, the physical and chemical data from cruises made from 1949-1959, inclusive, were subsequently published in Oceanic Observations of the Pacific (see footnote 1). The Data Reports for cruises prior to April 1953 contain parameter values for standard depths only. Subsequent Data Reports included parameter values for both observed and standard depths. MLR cruise numbers were dropped after 1958. In some cases, the Data Reports cover more than one cruise.

CCOFI         MLR         Period of Observation         SIO Ref.         Date of Report Report Processing           (5002)         11         Jan. 31-Feb. 19, 1950         none         Feb. 19, 1950           (5003)         12         Mar. 2-17, 1950         none         Mar. 28, 1950           (5004)         13         Apr. 4-17, 1950         none         May 21, 1950           (5005)         14         May 3-18, 1950         51-17         Aug. 15, 1950
(5003) 12 Mar. 2-17, 1950 none Mar. 28, 1 (5004) 13 Apr. 4-17, 1950 none May 21, 19
(5006) 15 June 6-23, 1950 51-19 Aug. 20, (5007) 16 July 6-24, 1950 51-28 Aug. 30, (5008) 17 Aug. 5-19, 1950 51-32 Dec. 26, (5009) 18 Sept. 6-22, 1950 51-37 Nov. 1, 19 (5010) 19 Oct. 11-20, 1950 51-29 Sept. 12, (5011) 20 Nov. 9-28, 1950 51-47 Nov. 15, 19 (5010) 21 Jan. 9-29, 1951 51-55 Dec. 15, (5102) 22 Feb. 6-25, 1951 52-6 Feb. 20, (5202) 34* Feb. 6-23, 1952 52-24 May 1, 199 (5203) 35 Mar. 5- Apr. 3, 1952 52-36 July 15, (5204) 36 April 2-28, 1952 52-37 July 30, 5206 (38) June 3-21, 1952 56-27 Oct. 22, 5207 (39) July 2-24, 1952 56-27 Oct. 22, 5207 (39) July 2-24, 1952 56-27 Oct. 22, 5207 (39) July 2-24, 1952 56-32 Oct. 29, 5208 (40) Aug. 8-19, 1952 57-2 Jan. 22, 5210 (42) Oct. 8-22, 1952 57-13 March 19, 5211 (43) Nov. 6-21, 1952 57-22 May 14, 1952 10 (42) Oct. 8-22, 1952 57-22 May 14, 19, 5210 (42) Oct. 8-26, 1953 57-32 Aug. 7, 16, 5302 (45) Feb. 4-26, 1953 57-33 Aug. 15, 5303 (46) Mar. 6-Apr. 5, 1953 57-33 Aug. 15, 5304 (47) Mar. 31-Apr. 19, 1953 58-13 Feb. 20, 5305 (48) Apr. 29-May 30, 1953 58-14 Mar. 1, 19, 5305 (48) Apr. 29-May 30, 1953 58-14 Mar. 1, 19, 5306 (49) June 3-July 3, 1953 58-14 Mar. 1, 19, 5306 (49) June 3-July 3, 1953 58-14 Mar. 1, 19, 5306 (49) June 3-July 3, 1953 58-14 Mar. 1, 19, 5306 (49) June 3-July 3, 1953 58-14 Mar. 1, 19, 5306 (50) July 8-27, 1953 58-23 Aug. 1, 18, 5306 (51) Aug. 5-30, 1953 58-24 Aug. 1, 18, 5306 (52) Sept. 9-19, 1953 58-23 Aug. 1, 18, 5308 (51) Aug. 5-30, 1953 58-44 Aug. 1, 18, 5311 (54) Nov. 9-14, 1953 58-44 Aug. 1, 18, 5312 (55) Dec. 1-13, 1953 58-44 Aug. 1, 18, 5400 (56) July 8-27, 1954 58-65 Sept. 15, 5400 (61) June 2-23, 1954 59-35 Jan. 26, 5408 (63) Aug. 18-Sept. 10, 1954 59-42 May 18, 19, 5400 (64) Sept. 18-Oct. 17, 1954 59-42 May 18, 19, 5400 (65) Oct. 6-24, 1954 59-43 June 1, 19, 5400 (66) Sept. 18-Oct. 17, 1954 59-43 June 1, 19, 5410 (66) Nov. 10-16, 1954 59-43 June 1, 19, 5410 (66) Nov. 10-16, 1954 59-43 June 1, 19, 5410 (66) Nov. 10-16, 1954 60-1 Sept. 10, 5412 (67) Nov. 30-Dec. 16, 1954 60-1 Sept. 10, 5412 (67) Nov. 30-Dec. 16, 1954 60-1

<sup>\*</sup> Physical and chemical data reports apparently were not issued for MLR cruises 23-33.

Cru	ise	Period of Observation	SIO Ref.	Data of Panant
CCOFI	MLR	reriod of observation	SIU REI.	Date of Report
5501	(58)	Jan. 13-29, 1955	59-36	Oct. 15, 1959
5502	(69)	Feb. 9-23, 1955	59-36	Oct. 15, 1959
5503	(70)	Mar. 8-22, 1955	59-37	May 1, 1959
5504	(71)	Apr. 5-22, 1955	59-37	May 1, 1959
<b>5</b> 505	(72)	May 12-June 7, 1955	59-44	May 18, 1959
5507	(74)	July 8-23, 1955	59-46	May 18, 1959
5509	(76)	Sept. 10-25, 1955	59-47	May 19, 1959
5510	(77)	Oct. 16-30, 1955	60-2	June 15, 1959
5511	(78)	Nov. 8-20, 1955	60-3	July 1, 1959
5512	(79)	Nov. 29-Dec. 16, 1955	60-4	July 15, 1959
5601	<del>-</del>	Jan. 5-18, 1956	60-5	Aug. 27, 1959
5602		Feb. 3-21, 1956	60-5	Aug. 27, 1959
5603	••	Mar. 4-19, 1956	60-6	Nov. 11, 1959
5604		Apr. 5-27, 1956	60-6	Nov. 11, 1959
5605	-	May 4-22, 1956	60-34	Apr. 20, 1960
5606	-	May 26-June 25, 1956	60-34	Apr. 20, 1960
5607		July 6-25, 1956	60-35	Apr. 27, 1960
5608 5609	-	Aug. 7-19, 1956 Sept. 5-17, 1956	60-35 60-35	Apr. 27, 1960
5610	-	Sept. 27-Oct. 5, 1956	60-35	Apr. 27, 1960 Apr. 27, 1960
5611	-	Oct. 30-Nov. 5, 1956	60-35	Apr. 27, 1960
5612	-	Nov. 24-Dec. 21, 1956	61-22	Apr. 24, 1961
5701	(92)	Jan. 4-19, 1957	58-22	Mar. 5, 1958
5702	(93)	Feb. 6-26, 1957	58-22	Mar. 5, 1958
<b>57</b> 03	(94)	Mar. 6-30, 1957	58-24	Aug. 1, 1958
5704	(95)	Apr. 5-30, 1957	58-24	Aug. 1, 1958
5705	(96)	May 9-23, 1957	58-33	July 15, 1958
5706	(97)	June 5-23, 1957	58-33	July 15, 1958
5707	(98)	July 8-Aug. 3, 1957	58-63	Aug. 20, 1958
5708	(99)	Aug. 9-27, 1957	58-63	Aug. 20, 1958
5709	(100)	Sept. 4-21, 1957	58-64	Sept. 15, 1958
5710	(101)	Oct. 4-Nov. 8, 1957	58-64	Sept. 15, 1958
5711	(102)	Nov. 16-25, 1957	58-64	Sept. 15, 1958
5712	(103)	Dec. 12-20, 1957	58-64	Sept. 15, 1958
5801	-	Jan. 8-Feb. 2, 1958	59-3 59-8	Jan. 15, 1959 Feb. 1, 1959
5802 5803	<u>-</u>	Feb. 7-24, 1958 Feb. 27-Mar. 21, 1958	59-8	Feb. 1, 1959
5804	(108)	Mar. 30-Apr. 27, 1958	59-14	Mar. 1, 1959
5805		May 3-22, 1958	59-28	Dec. 10, 1958
5806	` _ ′	June 4-26, 1958	59-29	Jan. 5, 1959
5807		June 30-July 22, 1958	59-33	Feb. 24, 1959
5808	(330)	Aug. 6-21, 1958	59-48	Mar. 11, 1959
5809	(112)	Sept. 4-19, 1958	58-84	Dec. 1, 1958
5810	-	Oct. 8-Nov. 6, 1958	59-49	Apr. 17, 1959
5811	-	Nov. 13-24, 1958	59-50	Apr. 22, 1959
5812		Dec. 1-11, 1958	59-50	Apr. 22, 1959
5901	-	Jan. 7-29, 1959	59-58	Sept. 17, 1959
5902	-	Feb. 6-28, 1959	60-13	Oct. 21, 1959
5903		Mar. 11-28, 1959	60-36	Feb. 15, 1960

Cruise		Period of Observation	SIO Ref.	Date of Report
CCOFI	MLR	Terrod or observation	310 Ke1.	bate of Report
5904	_	Apr. 7-26, 1959	60-37	June 29, 1960
5905	_	May 6-27, 1959	60-38	Apr. 26, 1960
5906	_	June 3-30, 1959	60-39	Apr. 15, 1960
5907	_	July 9-Aug. 3, 1959	60-40	July 6, 1960
5908	-	Aug. 13-31, 1959	61-19	Sept. 12, 1961
5909	-	Sept. 9-Oct. 1, 1959	61-19	Sept. 12, 1961
5910	-	Oct. 8-30, 1959	61-20	Apr. 24, 1961
5911	-	Nov. 16-25, 1959	61-21	Apr. 24, 1961
5912	-	Dec. 9-19, 1959	61-21	Apr. 24, 1961
6001	-	Jan. 8-Feb. 13, 1960	61-23	June 20, 1961
6002	-	Feb. 11-Mar. 3, 1960	62-3	Aug. 18, 1961
6003	-	Mar. 10-29, 1960	62-5	Aug. 18, 1961
6004	-	Mar. 29-Apr. 30, 1960	62-6	Aug. 10, 1961
6005	~	May 13-29, 1960	62-7 62-8	Aug. 18, 1961
6006 6007 <b>-</b> 8	<u>-</u>	June 14-30, 1960 July 12-Aug. 14, 1960	62-8 62-9	Aug. 18, 1961 May 1, 1962
6008	_	Aug. 10-22, 1960	62-10	May 17, 1962
6009	-	Sept. 9-21, 1960	62-10	May 17, 1962
6009-10	_	Sept. 22-0ct. 22, 1960	62-10	May 17, 1962
6101-2	_	Jan. 5-Feb. 20, 1961	61-24	June 29, 1961
6103	-	Mar. 8-13, 1961	62-15	Dec. 28, 1961
6104-5	-	Apr. 4-May 12, 1961	62-15	Dec. 28, 1961
6105	-	May 17-29, 1961	62-15	Dec. 28, 1961
6107-8	-	June 27-Aug. 1, 1961	62-16	July 26, 1962
6108	-	Aug. 13-28, 1961	62-16	July 26, 1962
6109	-	Aug. 31-Sept. 19, 1961	62-17	July 26, 1962
6110-11	-	Oct. 10-Nov. 12, 1961	62-17	July 26, 1962
6201-2	-	Jan. 12-Feb. 15, 1962	62-26	Nov. 19, 1962
6203-4	-	Mar. 15-May 2, 1962	63-9	Feb. 15, 1963
6207-8 6210-11	-	July 10-Aug. 29, 1962	62 <b>-</b> 23	Oct. 4, 1962
6212	_	Oct. 5-Nov. 18, 1962 Dec. 17-19, 1962	63-25 63-25	Aug. 22, 1963 Aug. 22, 1963
6301-2	-	Jan. 10-Feb. 23, 1963	64-2	Mar. 5, 1963
6304	•••	Apr. 9-May 24, 1963	64-13	Mar. 2, 1964
6306	_	June 25-26, 1963	64-13	Mar. 2, 1964
6307	-	July 10-Aug. 8, 1963	64-18	May 25, 1964
6309	-	Sept. 3-29, 1963	64-18	May 25, 1964
6310		Oct. 2-29, 1963	65-1	Oct. 9, 1964
6311		Nov. 28-29, 1963	65-1	Oct. 9, 1964
6311 (E1 Go	olfo)	Nov. 9-Dec. 7, 1963	65-1	Oct. 9, 1964
6401	•••	Jan. 9-Mar. 4, 1964	65-7	Apr. 5, 1965
6404	-	Apr. 10-May 1, 1964	66-20	Sept. 1, 1966
6407 6410	_	June 15-Aug. 4, 1964 Oct. 2-Nov. 1, 1964	66-20 65-18	Sept. 1, 1966
6501	_	Jan. 6-Feb. 18, 1965	66-4	Oct. 5, 1965 Dec. 9, 1965
6504	_	Mar. 31-Apr. 24, 1965	67 <b>-</b> 16	(undated)
6505 (E1 Go	olfo II)	May 14-June 17, 1965	67-16	(undated)
6507	-	June 15-Aug. 11, 1965	67-17	(undated)
6509	-	Aug. 31-Sept. 25, 1965	67-17	(undated)
				•

Cruise	s	Period of Observation	SIO Ref.	Date of Report
CCOFI	MLR	reriod or observation	310 NC1.	bate of Report
6601		Jan. 12-Feb. 7, 1966	68-3	(undated)
6602	_	Feb. 15-Mar. 6, 1966	68-3	(undated)
6604	_	Mar. 26-May 3, 1966	68-3	(undated)
6605		May 5-29, 1966	68-3	(undated)
6606		June 12-July 1, 1966	68-3	(undated)
6607	-	July 8-29, 1966	68-21	(undated)
6608	-	Aug. 5-25, 1966	68-21	(undated)
6609	-	Sept. 7-24, 1966	68-21	(undated)
6610	-	Oct. 8-27, 1966	69-2	(undated)
6611	-	Nov. 10-13, 1966	69-2	(undated)
6612	-	Dec. 2-19, 1966	69-2	(undated)
6707	-	June 19-July 20, 1967	69-8	(undated)
6712	-	Dec. 5-20, 1967	69-8	(undated)
6801	-	Jan. 7-26, 1968	71-3	(undated)
6804	-	Apr. 23-May 6, 1968	71-3	(undated)
<b>6</b> 806	-	May 31-June 22, 1968	71-3	(undated)
6901	-	Jan. 7-30, 1969	76-14	Sept. 1, 1976
6902	-	Jan. 26-Mar. 11, 1969	76-14	Sept. 1, 1976
6904	-	Apr. 2-26, 1969	77-22	Nov. 1, 1977
6905	-	May 5-29, 1969	77-22	Nov. 1, 1977
6906	-	June 9-28, 1969	77-22	Nov. 1, 1977
6907	-	July 10-29, 1969	79 <b>-</b> 7	May 15, 1979
6908	-	Aug. 6-Sept. 8, 1969	79-7	May 15, 1979
6909	-	Sept. 11-Oct. 7, 1969	79 <b>-</b> 7	May 15, 1979
6910	-	Oct. 9-Nov. 8, 1969	79-29	Feb. 15, 1980
6912	-	Nov. 13-Dec. 17, 1969	79-29	Feb. 15, 1980
7008	-	Aug. 17-Oct. 2, 1970	79-30	Feb. 15, 1980
7102	-	Feb. 8-Apr. 5, 1971	79-30	Feb. 15, 1980
7201	-	Jan. 3-26, 1972	80-21	Oct. 15, 1980
7202	-	Feb. 1-27, 1972	80-21	Oct. 15, 1980

Appendix 2: The Packed Binary Format for the CalCOFI Oceanographic Data File

The packed binary format for data from Nansen casts or STD or CTD data obtained at CalCOFI stations was designed primarily to reduce the size of the file stored on magnetic tape. This was accomplished in part by use of the binary format and in part by use of variable length records. Each record contains identification fields, sea state and surface weather observations, measured and computed parameter values at both observed and standard depths. The number of depths varies considerably from record to record, which accounts for the variation in record length. To convert the file to fixed length records with no other change in format would require a record size of 300 (48 bit) words. Most of the records, however, are less than half this size.

The first six words (36 char) of each record contain location-time identification fields and ship code, all in ASCII characters. The next three words (18 char = 144 bits) contain surface observations along with depth to the bottom, depth of the cast and the total number of four word (observed depth) groups and three word (standard depth) groups to follow in the record. These groups are ordered by increasing depth and are intermixed throughout the record. A four-bit field at the right hand end (least significant bits) of the first word in each observed or standard depth group contains a type code. Code values of 3 or 4 indicate observed depths and 6 or 7 indicate standard depths.

The location of record elements in the following list are given in terms of a consecutive word number within each group and the bit field location within each word. The bits are numbered 0 to 47 beginning with the right-most bit of a word; however, the elements are accessed from left to right (high order to low order bits) in each word.

#### --- IDENTIFICATION GROUP: 36 CHAR ---

Element	Form	Char. Field
CalCOFI Station	LLLSSS	1-6
Year-Month-Day Code	YYMMDD	7-12
Latitude-Deg. and min. (decimal point implied; H="N" or "S")	DDMM.MH	13-18
Longitude-Deg. and Min. (decimal point implied; H = "E" or "W")	DDDMM.MH	19-25
Marsden Square	MSQ	26-28
Hour (decimal point implied)	нн.н	29-31
Ship Code	SS	32-33
Blank	bbb	34-36

# --- SURFACE OBSERVATIONS GROUP: 3 WORDS ---

Element	Word	Bit-Field
Water Color	1	47-41
Water Transparency	1 1	40-34
Wave Direction - Deg.		33-27
Wave Height Flag	1	26
1 = wave height unknown		
Sea State or Wave Height Code	1	25
1 = sea state		
2 = wave height		
Wave Height-Meters (Sea State)	1	24-21
Wave Period Flag	1	20
1 = wave period unknown		
Wave Period - Sec.	1	19-16
Wind Direction - Deg.	1	15-9
Wind Speed - Beaufort before 1969,	1	8-2
knots since 1969		
Zero Fill	1 2	1-0
Depth to Bottom - Fathoms	2	47-33
before 1969, meters since 1969		
Depth of Cast - hundreds of meters	2	32-26
No. of Obs and STD Depths	2	25-18
Barom. Pressure - Millibars	2	17-6
Zero Fill	2	5-0
Dry Bulb Temperature - °C	3	47-37
Wet Bulb Temperature - °C	3	36-26
Cloud Type Code	3	25-21
Cloud Amount Code	3	20-16
Visibility - Naut. Mi.	3	15-11
Weather Code	2 2 2 3 3 3 3 3 3	10-3
Zero Fill	3	2-0

# --- OBSERVED DEPTH DATA GROUP: 4 WORDS ---

Element	Word	Bit Field
Depth Reliability Code 1 = doubtful	1	47
Observed Depth - Meters	1	46-34
Sigma-t - g/L	1	33-19
Sound Velocity - M/Sec.	1	18-4
Observed Depth Code - 3 or 4	1	3-0
Temperature Reliability Code 1 = doubtful	2	47
Sea Water Temperature - °C	2	46-30
Salinity Reliability Code - 1 = doubtful	2	29
Sea Water Salinity - Oxygen Reliability Code -	2	28-12

<pre>1 = doubtful</pre>		11
Dissolved Oxygen - ml/L	2	10-0
Phosphate - Phosphorus - $\mu g$ at/L	3	47-32
Total Phorphorus - $\mu g$ at/L	3	31-16
рН	3	15-0
Nitrite - Nitrogen - $\mu$ g at/L	4	47-32
Nitrate - Nitrogen - μg at/L	4	31-16
Silicate - Silicon - μg at/L	4	15-0

# --- STANDARD DEPTH DATA GROUP: 3 WORDS ---

<u>Element</u>	Word	Bit Field
Depth Reliability Code - 1 = doubtful	1	47
Standard Depth - Meters	1	46-34
Sigma-T g/L	1	33-19
Sound Velocity - M/Sec.	1	18-4
Standard Depth Code - 6 or 7	1	3-0
Temperature Reliability Code - 1 = doubtful	2	47
Sea Water Temperature - °C	2	46-30
Salinity Reliability Code - 1 = doubtful	2	29
Sea Water Salinity - o/oo	2 2	28-12
Oxygen Reliability Code - 1 = doubtful	2	11
Dissolved Oxygen - ml/L	2	10-0
Thermosteric Anomaly - cl/ton	3	47-26
Geopotential Anomaly - Dyn. Meters	2 3 3 3	25-10
Zero Fill	3	9-0

There are no specific codes to identify missing data in the packed binary file. Whenever a record element was missing in the original source file, the associated bit field in the binary file was zero-filled. Zero values can be interpreted as missing data for most of the important elements without ambiguity.

#### APPENDIX 3: Deck Set-Up for CCFIHYDRO

The general assemblage of control and data cards needed to run the program on the Burroughs B7800 is as follows:

- ? BEGIN JOB
- ? RUN SERVICE/OPINFO TAPENAME, 9/800, RING IN, PURGE, OU
- ? RUN CCFIHYDRO
- ? FILE CCFI (FILE ATTRIBUTES)
- ? FILE SUBFIL (FILE ATTRIBUTES)
- ? FILE OCNDAT (FILE ATTRIBUTES)
- ? FILE CALTAP (TITLE = TAPENAME)
- ? DATA

RUN SPECIFICATION CARD DATA SELECTION CARDS

? END JOB

The SERVICE/OPINFO cards are needed only if tape output (FILE CALTAP) is required. CCFI is the input file in packed binary format. Only one of the three following file cards are needed, depending on the output desired. The format and entries for run specification and data selection cards are described in the main body of this document.

The following example illustrates the kind of retrieval application for which the program was designed. The object will be to extract data at standard depths from 0 to 200 meters at stations on Line 90 out to station 100, and to separate the data by month and store it in 12 working files in the simple binary format.

A direct approach to this task is to set up the control card specifications for data selection mode #7 (see Table 3) and prepare one set of data selection cards for the standard depths, a second set for the Line 90 stations and 12 sets of cruise cards, one for each month. Then the job could be done with 12 runs of CCIFHYDRO, one for each month. The task can be performed much more efficiently, however, by making a preliminary run to create a SUBFIL which will include only the desired stations.

The job assembly for the preliminary run is listed below:

- ? BEGIN JOB
- ? RUN CCFIHYDRO
- ? FILE CCFI (TITLE = CCOFI/PAKDAT)
- ? FILE SUBFIL (TITLE = SUBFIL/LINE90)
- ? DATA

(RUN SPECIFICATION CARD-b means space)

blineb90bsubfilbbbbbbbbbbbbsubfilbstationsballb#

(1st set of data selection cards - stations)

bbbbb11bb090028bb090030bb090032b b090037bb090045bb090053bb090060b b090070bb090080bb090090bb090100b

#### ? END JOB

Fields 5 and 6 of the specification card are not needed for this run. Fields on data selection cards should be entered continuously in cols. 1 through 80. In the example above, this would require all of one card and the first 16 cols. of a second card. The station list includes 11 stations. The small letter "b" in the example indicates a blank space. The output file, SUBFIL/LINE90 is much smaller than the source file, CCOFI/PAKDAT, and will be used as the input file for the remaining 12 runs. The job assembly listed below is for the January data. That for other months would be similar.

- ? BEGIN JOB
- ? RUN CCFIHYDRO
- ? FILE CCFI (TITLE = SUBFIL/LINE90)
- ? FILE OCNDAT (TITLE = OCNDAT/LINE90/JAN)
- ? DATA

(RUN SPECIFICATION CARD)

blineb90bJANbCRUISESbbbbDEPTHbDISTNbSTDb1b#

(1 set of data selection cards - depths)

bbbbb10bbbb000bbbb010bbbb020b bbbb030bbbbb050bbbbb075bbbbb100b bbbb125bbbbb150bbbbb200

(2nd set of data selection cards - stations. These will be the same as the 1st set for the preliminary run, shown above)

(3rd set of data selection cards - cruises. Only a few cruises are included for brevity)

bbbbb08bbbb6001bbbb6101bbbb6201b bbb6301bbbb6401bbbb6501bbbb6601b bbbb6701

The output file, OCNDAT/LINE 90/JAN, resulting from this run will contain data from 8 January cruises for 11 stations on Line 90.

# Appendix 4: Data Extraction from the Packed Binary Oceanographic File With Procedure UNPACKV

UNPACKV is an ALGOL procedure designed to unpack bit-fields and extract data values from records read from the packed binary data file. It must be bound or compiled into a host program and the host program must contain an input routine to read records from the source file into an array which will be passed to the procedure as a parameter. UNPACKV has seven parameters, described below in the order that they must be entered in the procedure statement.

VRBL Integer code number of the data parameter to be extracted--see parameter list.

SAMTYPE Character string denoting the group in the source record from which the sample is to be extracted. Valid entries are:

"STAT" for parameters in the surface observations;

"OBS" for data at observed depths only;

"STD" for data at standard depths only;

"OBSTD" for data at both observed and standard depths.

MINDPTH Integer value specifying minimum depth for extracting data.

MAXDPTH Integer value specifying the maximum depth for extracting data.

REC Name of an INTEGER array into which a record from the source file has been read.

NUMWORDS Integer variable which has been assigned the length of the record in 48-bit words.

VRBLIST Name of a REAL array into which the data values extracted by UNPACKV will be stored.

DPTHLIST Name of a REAL array into which the depths of the extracted data values will be stored by UNPACKV. The elements of this array will correspond one for one with those of VRBLIST.

NUMDPTHS Integer variable returned by UNPACKV with the number of depths for which data were extracted.

The data parameters corresponding to the code values of VRBL are listed below. Note that only certain SAMTYPE entries are associated with each code value. If an invalid combination is submitted, UNPACKV will return a value of -1 for NUMDPTHS and no data will be extracted. If "STAT" is entered for SAMTYPE then the procedure will return 1 for NUMDPTHS and 0 in the first element of DPTHLIST.

VRBL	SAMTYPE	SAMPLE RETURNED IN VRBLIST
1	STAT	Water - 4 digits, 1st-2nd: Transparency
_	A#4#	3rd-4th: Color
2	STAT	Waves - 7 digits 1st: if=1, 5th digit is sea state,
		otherwise wave height
		2nd: if=1, period unknown
		3rd: if=1. height unknown
		4th: period
		5th: sea state or wave height
•	CTAT	6th-7th: direction.
3	STAT	Wind - 4 digits, 1st-2nd: direction
		3rd-4th: speed in Beaufort (before
		1969) or knots (since 1969)
4	STAT	Depth to bottom - fathoms
		(before 1969) or meters
_	0717	(since 1969).
5 6	STAT STAT	Depth of cast. Number of depths
O	SIAI	(OBS + STD)
7	STAT	Barometric pressure.
8	STAT	Air temperature - dry bulb.
9	STAT	Air temperature - wet bulb.
10	STAT	Clouds - 3 digits,
		1st: if=1, type unknown 2nd: amount
		3rd: type.
11	STAT	Visibility
12	STAT	Weather - 3 digits,
		1st: if=1, WMO code 4501 used
		if=0, WMO code 4677 used
10	000 000 000	2nd-3rd: weather code
13 14	OBS, STD or OBST OBS, STD or OBST	D Sigma-t D Sound velocity
15	OBS, STD or OBST	
•	,	3 or 4 = observed depth,
		6 or 7 = standard depth.
16	OBS, STD or OBST	
17	OBS, STD or OBST	
18 19	OBS, STD or OBST OBS	D Oxygen Phosphorous - phosphate
20	OBS	Total phosphorus
21	OBS	рН
22	OBS	Nitrite - nitrogen
23	OBS	Nitrate - nitrogen
24	OBS	Silicate - silicon
25 26	STD STD	Thermosteric anomaly Geopotential anomaly
20	310	Geopo central anomary

UNPACKV does not extract fields from the identification group of a packed binary record. These fields are all character strings (see Appendix 2) and

can be obtained from the array into which the record was read, as will be shown in the example to follow, which is based on the input routine from CCFIHYDRO.

### **EXAMPLE**

```
BEGIN COMMENT READ PACKED BINARY FILE:
  FILE CCFI (KIND=1, MAXRECSIZE = 20, BLOCKSIZE = 60,
     AREASIZE = 60, FILETYPE = 6);
  ARRAY REC [0:300], DPTHLIST [0:50], VRBLIST [0:26, 0:50];
  INTEGER VRBL, MINDPTH, MAXDPTH, NUMWORDS, NUMDPTHS, STNCODE,
     YYMMDD, HR, MSQ;
  REAL SAMTYPE, LATD, LATM, NS, LOND, LONM, EW, SHIP;
  BOOLEAN READIT;
  LABEL NXTREC, EOF;
NXTREC:
  READIT:= READ (CCFI, 300, REC);
 COMMENT THIS STATEMENT READS A RECORD INTO ARRAY REC AND STORES
     THE VALUE OF THE I/O DESCRIPTOR IN READIT. THE I/O DESCRIPTOR
     CONTAINS THE STATUS OF THE I/O ACTION, INCLUDING THE LENGTH OF
     THE RECORD AND WHETHER AN END OF FILE WAS REACHED;
  IF REAL (READIT.[9:1]) = 1 THEN GO TO EOF;
  NUMWORDS: = REAL (READIT.[47:20]):
  COMMENT THIS IS THE RECORD LENGTH;
  READ (REC, <216, 12, 13, A1, 213, A1, 13, 12, A2>, STNCODE,
     YYMMDD, LATD, LATM, NS, LOND, LONM, EW, MSQ, HR, SHIP);
  COMMENT THIS ASSIGNS THE ID FIELDS TO INTEGER OR REAL VARIABLES.
     THEN MINUTES OF LAT AND LON ARE CHANGED TO DEGREES AND THE DECIMAL
     PLACES RESTORED WHERE APPROPRIATE:
  LATD: = LATD + LATM/600;
  LOND: = LOND + LONM/600; HR: = HR/10;
  MINDPTH: = 0; MAXDPTH: = 500;
  SAMTYPE: = "STAT";
  FOR VRBL: = 1 STEP 1 UNTIL 12 DO
  UNPACKV (VRBL,SAMTYPE, O. O. REC, NUMWORDS, VRBLIST [VRBL, *],
     DPTHLIST, NUMDPTHS);
  COMMENT THE FIRST CALL TO UNPACKY GETS THE ID FIELDS AND SURFACE
     OBSERVATIONS:
  SAMTYPE: = "OBSTD";
  FOR VRBL: = 13 STEP 1 UNTIL 18 DO
  UNPACKV (VRBL, SAMTYPE, MINDPTH, MAXDTH, REC, NUMWORDS,
    VRBLIST [VRBL,*], DPTHLIST, NUMDPTHS);
  COMMENT THE NEXT CALL GETS THE DATA PARAMETERS THAT ARE IN BOTH
     OBS DEPTH AND STD DEPTH GROUPS:
  SAMTYPE: = "OBS":
  FOR VRBL: = 19 STEP 1 UNTIL 24 DO
  UNPACKY (VRBL, SAMTYPE, MINDPTH, MAXDPTH, REC, NUMWORDS,
     VRBLIST[VRBL,*], DPTHLIST, NUMDPTHS);
  COMMENT THE THIRD CALL GETS DATA PARAMETERS AT OBS DEPTHS ONLY:
  SAMTYPE: = "STD":
  FOR VRBL: = 25, 26 DO
  UNPACKY (VRBL, SAMTYPE, MINDPTH, MAXDPTH, REC, NUMWORDS,
     VRBLIST [VRBL,*], DPTHLIST, NUMDPTHS);
```

COMMENT NOW ALL 26 FIELDS ARE STORED IN ARRAY VRBLIST. THIS DATA MUST BE PROCESSED BEFORE READING THE NEXT RECORD;
GO TO NXTREC;
EOF: END.

# Appendix 5: Utility Programs for OCNDAT Files

The OCNDAT file format has fields for the six data parameters which occur most frequently in the CalCOFI oceanographic data file: temperature, salinity, oxygen, sigma-t, thermosteric anomaly and geopotential anomaly. It provides also for the identification parameters: station code, depth, year, month, day and cruise code. It excludes all other identification fields, all of the surface observations and the nutrients.

Logical records in OCNDAT files consist of nine 32-bit fields for a record length of 36 bytes. The physical record size (BLOCKSIZE) is 40 logical records, or 1440 bytes. For applications which do not require the excluded elements, the OCNDAT format is more efficient and easier to use than the packed binary format described in Appendix 2.

OCNDAT files are composed of one or more data sets, each of which begins with two header records. The first header record has a character string field of 28 bytes, containing file identification, followed by 2 integer fields of 32 bits each. The latter contain a data set sequence number and the number of data records in the data set, respectively. The second header record contains a single character string field of 36 bytes which identifies the data set. Data records each contain nine integer fields as described below:

- 1st. Six-digit CalCOFI station code or four-digit cruise code.
- 2nd. Six-digit year-month-day code.
- 3rd. Depth (meters).
- 4th. Temperature \* 1000 (°C).
- 5th. Salinity \* 1000 (o/oo).
- 6th. Dissolved oxygen \* 1000 (ml/L).
- 7th. Sigma-t \* 1000 (g/L).
- 8th. Thermosteric anomaly \* 100 (cl/Ton).
- 9th. Geopotential anomaly \* 10000 (Dyn. Meters).

The high order bit in each 32-bit field of an OCNDAT data record is sign bit. Negative values are represented by two's-complement notation.

Several utility programs have been written to manipulate OCNDAT files. They are described in the following sections.

#### PRTOCNDAT

This program copies an OCNDAT file to an EBCDIC (or ASCII) disk file or to an EBCDIC (or ASCII) tape file and/or prints it. The file names and attributes declared in the program for disk and tape are:

FILE OCNEBC (KIND=DISK, MAXRECSIZE=72, BLOCKSIZE=1440, UNITS= CHARACTERS, AREASIZE=30, FLEXIBLE).

FILE OCNTAP (KIND=TAPE9, MAXRECSIZE=72, BLOCKSIZE=1440, UNITS=

CHARACTERS. EXTMODE=EBCDIC. LABELTYPE=OMITTEDEOF).

The two header records for each data set in the OCNDAT file are written as a single header record in the output file with the format: (4A6, C4, 2I4, 6A6). Data records are written as 8 byte fields with the format: (3I8, 4F8.3, F8.2, F8.4).

A control card must be submitted at run time with one or more of the output options "PRINT", "DISK" and/or "TAPE." The format for the control card is (3A6).

The option(s) may be entered in any order but must be right justified in each field. If "TAPE" is one of the options submitted, the program will copy 1 or more OCNDAT files into a single tape file. The TITLES of each file to be copied to tape must be entered on one or more subsequent data cards in the following format (I6,X1,3(4A6),X1/(X7,3(4A6),X1)) where the first field (I6) must contain the number of files (right justified) to be copied and the respective TITLES, up to 23 characters in length, should begin in columns 8, 32 or 56 and each one terminate with a period. The program output will list the name of every file copied. The deck set-up is shown in the following examples.

## Example 1:

?BEGIN JOB ?RUN PRTOCNDAT ?FILE OCNDAT (TITLE=OCNDAT/LINE 90/JAN) ?DATA bPRINTbbDISK ?END JOB

# Example 2:

?BEGIN JOB ?RUN PRTOCNDAT ?DATA bbTAPE bbbbb2b0CNDAT/LINE90/JAN.bbbbbb0CNDAT/LINE90/FEB. ?END JOB

The 1st example prints the file named in the FILE card and also copies it to a new, EBCDIC disk file. The 2nd example copies two files to a single file on tape. No FILE card is needed because the files copied are named in the data cards.

#### 2. MAKOCNSUB

This program copies selected records from an OCNDAT file to a new file in the same format. The selection criteria are either station codes, cruise codes or standard depth codes. The internal name of the output file is OCNSUB; it is declared with the same attributes as OCNDAT. The selection parameter along with others must be submitted on a control card at run time. Valid entries for the selection parameter are "STATIONS", "CRUISES" and "STDbDEPTH." It should be entered, left justified, in the first field (cols.

1-12) of the control card. Entries in all subsequent fields must be right justified, when applicable. If the selection parameter is "STDbDEPTH", the second field (cols. 13-18) must contain a valid standard depth code. Otherwise it is left blank. The third field (cols. 19-24) is for a print option and will cause all the data records, copied to the new file, to be printed if "PRINT" is entered. The next four fields have 6 characters each (cols. 25-48) and will be copied into the first 4 words of each header record of the new file, for identification. However, if OCNSUB is going to be used as a source file for the contouring program EDMAP2, the 4th and 5th fields will be interpreted by that program as the y and x coordinates, respectively, of a rectangular grid to which the data from OCNSUB will be fitted. In this case, valid entries for the 4th field (cols. 25-30) are "bDEPTH", "bDISTN" and "bDISTP", which will be interpreted as depths, station numbers and line numbers. Similarly, valid entries for the 5th field (cols. 31-36) are "bDISTN", "bDISTP" "bbTIME", which will be interpreted as station numbers, line numbers and months. The 6th and 7th fields are optional in any case. If "STATIONS" or "CRUISES" is entered as the selection parameter, a station list or cruise list must be submitted, following the control card. The format is (10(X1, I6, X1)) in which the first field must contain an integer specifying the number of station codes or cruise codes to follow. The list may take one or more cards. All fields are right justified. The deck set-up for running MAKOCNSUB is shown in the following examples.

# Example 1:

#### Example 2:

?BEGIN JOB
?RUN MAKOCNSUB
?FILE OCNDAT(TITLE=OCNDAT/ALLSTNS/JAN)
?FILE OCNSUB(TITLE=OCNDAT/10METERS/JAN)
? DATA
 (Run control card)
STDbDEPTHbbbbbb010bPRINTbDISTPbDISTNbSTDb10MbJAN
?END JOB

In the 1st example, data for 6 stations on Line 90 will be extracted from an OCNDAT file containing all January stations and copied to a new file. The data records will not be printed. In the 2nd example, data at the 10 meter depth will be extracted for all stations from the same file. In this case the selection criteria is specified in the second field of the control card, and no other data cards are needed.

#### ADDSTN

This program merges two OCNDAT files and creates a new file in the same format. OCNDAT files are normally structured such that each data set in the file contains a number of stations, sequentially ordered by ascending station codes, for one cruise. There are a varying number of records for each station, ordered by increasing depth. The data sets are ordered by ascending cruise code within the file. The program designates the two existing files as OLDFIL and ADDFIL, and the new file as NEWFIL. The procedure, essentially, is to copy records from OLDFIL to NEWFIL and insert records from ADDFIL when appropriate. If ADDFIL contains a cruise (data set) not in OLDFIL, the entire cruise is inserted. If the same cruise is in both OLDFIL and ADDFIL, then each station for that cruise found in ADDFIL and not found in OLDFIL is inserted in NEWFIL. When the same station is found in both files, only the one from OLDFIL will be copied. The program requires one data card to be submitted at run time with the format (4Å6), containing a character string which will be inserted into header records on NEWFIL for data set identification. Data set sequence numbers and record counts, will be tallied and entered in NEWFIL header records by the program. The deck set-up for running ADDSTN is shown in the following example.

#### Example:

?BEGIN JOB ?RUN ADDSTN ?FILE OLDFIL (TITLE=OCNDAT/102STNS) ?FILE ADDSTNS (TITLE=OCNDAT/36STNS) ?FILE NEWFIL (TITLE=OCNDAT/138STNS) ?DATA bDEPTHbDISTNb138STNSbALL ?END JOB

In this example records for 36 stations in ADDFIL will be merged, cruise by cruise, with records for 102 stations in OLDFIL to create a NEWFIL of 138 stations. The printer output will include all header records read from OLDFIL and ADDFIL and written to NEWFIL.

#### 4. SORTOCNDAT

This program sorts the records in an OCNDAT file using station code as a sorting key and creates a new file in the same format. The new file will consist of a single data set in which the records are ordered by ascending station codes. Cruise codes contained in the header records of the original file will not be in the new file. Up to 16000 stations can be sorted. The internal name of the sorted file is OCNSEQ. No control or data cards are required at run time. SORTOCNDAT is a prerequisite to MEANOCNSEQ, which is described in the next section. The deck set-up for running the program is shown in the following example.

#### Example:

?BEGIN JOB
?RUN SORTOCNDAT
?FILE OCNDAT (TITLE=OCNDAT/ALLSTNS/JAN)

?FILE OCNSEQ (TITLE=OCNSEQ/ALLSTNS/JAN) ?END JOB

The printer output will include the header records for each data set in the input file and a count of stations and records sorted.

### 5. MEANOCNSEQ

This program computes means and standard deviations at standard depths for each station on OCNSEQ files created by SORTOCNDAT. The outputs are stored in three new files, OCNMEA, OCNDEV and OCNCNT created in the OCNDAT format. The 4th through 9th fields of data records in OCNDAT files contain values of temperature, salinity, oxygen, sigma-t, thermosteric anomaly and geopotential anomaly. The corresponding fields in the new files will contain the means, standard deviations and counts of those parameters. The computation is straightforward for all of the data parameters except geopotential anomaly (9th field). In the OCNDAT files, the value stored in this field in any record was obtained by integration of specific volume anomaly from the surface to the depth represented by that record. MEANOCNSEQ computes the mean geopotential anomaly, for a station, at each standard depth from 0 to 500 meters. Then, the mean value at each depth is replaced by its difference from the mean value at 500 meters. The reference level for the mean values or geopotential anomaly is therefore changed from 0 to 500 meters, so that transports in upper layers, relative to that at 500 meters, can be computed from gradients between adjacent stations. For inshore stations which do not go to 500 meters, provision is made for supplying the requisite 500 meter reference values via data cards at run time. Determination of such values can be made by extrapolation from deeper stations farther offshore. MEANOCNSEQ will do the extrapolation automatically for all stations which do not reach 500 meters if the option AUTO is entered on a control card at run time. The procedure involves the two nearest stations outward from the shallow station on the same CalCOFI line. The increment of geopotential anomaly between the maximum depth of the shallow station and 500 meters is determined by extrapolation from the two deeper stations.

The program does not compute standard deviations for geopotential anomaly. A control card must be submitted at run time to designate options for printer output and the 500 meter reference level values. The control card format is: (2A6). If "bPRINT" is entered in the 1st field (cols. 1-6), the three output files will be printed. Otherwise they will not. If "bbAUTO" is entered in the 2nd field, the 500 meter reference values for shallow stations will be computed automatically. Otherwise, they must be submitted in one or more data card(s) following the control card. The format is:(X16,8R8.0) where the 1st field must contain the number of stations for which reference values are entered. The station codes and associated reference values are entered in subsequent pairs of field.

The deck set-up for MEANOCNSEQ is shown in the following example.

#### Example:

?BEGIN JOB
?RUN MEANOCNSEQ
?FILE OCNSEQ (TITLE=OCNSEQ/ALLSTNS/JAN)

In this example, the output files will be printed and 500 meter reference level data are given for two stations, 82047 and 83051. The portion of the card preceding the first field (cols. 1-16) contains identification for visual reference only.

#### 6. INTERPSTNS

This program performs horizontal interpolation from irregularly spaced stations on CalCOFI lines to regularly spaced points of a grid, superimposed on the CalCOFI station pattern. The interpolations are made only for points on the inshore side of station 60, on each line. The purpose of the program is to prepare source data files for the contouring program EDMAP2, so as to avoid introducing bias in horizontal gradients of the gridded data fields produced by that program. INTERPSTNS reads data from file OCNMEA, which can be any file with the OCNDAT format, and creates a new file, OCNSTN. Control cards submitted at run time designate whether the output file will be printed, and specify the points on the CalCOFI lines to which the interpolations will be made. The first control card has the format: (A6), and will invoke the print option if the entry is "bPRINT". Subsequent control cards have the format: (10R8.0). The first field (cols. 1-8) must contain the interval, specified in station number units, for the interpolated values. The second field must contain the number of CalCOFI lines in the input file. Subsequent pairs of fields will contain line numbers followed by station numbers which designate the closest inshore points to which values will be interpolated. A designated point may be inshore of the first station on a line, in which case extrapolation will be used. The deck set-up is shown in the following example.

# Example:

For this example, it is assumed that the input file contains data for only three lines. If a station is encountered which is not on one of the lines designated in the control cards, the program will terminate. The 1st control card specifies that the output file will be printed. The 2nd one

specifies that interpolations will be made at intervals of 2.5 station units beginning with station 52.5 on line 80, 27.5 on line 90 and 30.0 on line 100. If a point for interpolation is coincident with an actual station, the data is simply copied from the input file to the output file.

#### 7. INTERPSIGT

This program computes the depth of a designated value of sigma-t at each station in an OCNDAT file and determines the values of data parameters at that depth. Salinity and oxygen are computed by linear interpolation from standard depths. Temperature is computed from the designated sigma-t value and the interpolated salinity. Acceleration potential is computed from the formula:

Acc.Pot.=
$$(GP)+D*(SPV(S,T,D)-SPV(35,0,D))$$

where D is the depth of the designated sigma-t surface, (GP) is the interpolated value of geopotential anomaly at D, SPV(S,T,D) is the specific volume at the interpolated salinity and temperature at depth D, and SPV (35,0,D) is the specific volume at standard values of salinity and temperature, at depth D. These values are used to create a new file, OCNSIG, in the same format as the input file, OCNSTN, except that the depth of the designated sigma-t replaces sigma-t in the 7th field and acceleration potential replaces geopotential anomaly in the 9th field, and there will be only one record for each station. A control card must be submitted at run time with the designated value of sigma-t and a 12-character string which will be inserted in header records of the output file for identification. The control card format is: (R6.0,2A6), in which the first field must contain the sigma-t value and the two alpha fields must contain the character string. The deck set-up is shown in the following example.

#### Example:

?BEGIN JOB ?RUN INTERPSIGT ?FILE OCNSTN (TITLE=OCNSTN/LINES80T0100) ?FILE OCNSIG (TITLE=OCNSIG/LINES80T0100) ?DATA (control card) bb25.8bSIGT258bJan ?END JOB

This example uses the output of INTERPSTNS for input and computes values of data parameters at depths where sigma-t = 25.8.

## 8. FIT/HARMONICS

This program computes annual and semi-annual harmonic coefficients for data parameters in OCNDAT files by the method of least squares. The coefficients are related to the data parameters by the formula:

Y = A1\*COS(X)+B1\*SIN(X)+A2\*COS(2X)+B2\*SIN(2X)+C

where Y is the data parameter, X is the time coordinate on an annual cycle,

and Al. Bl. A2. B2 and C are the coefficients. In computing the latter, X is determined from the year-month-day code in the second field of OCNDAT records. The input file will generally contain data from all available cruises for one or more stations. It must have been created by SORTOCNDAT, which sorts OCNDAT files by station code and combines data sets. Coefficients are computed for each of the six OCNDAT data parameters at each standard depth to 600 meters for each station, provided the sufficiency criterion is met. This requires that a station must have been occupied at least once in each of the six periods: Jan. 1-Feb. 28, Mar. 1-Apr. 28, Apr. 29-June 26, June 27-Aug. 24, Aug. 25-Oct. 22, Oct. 23-Dec. 20. The period Dec. 21-31 is a time of little or no CalCOFI activity. The above criterion is based on the premise that a least squares fit for five coefficients can be computed with only six data values provided they are reasonably spaced through the year. The output of the program will be stored in file HMCOEF in which each record has a length of 32 (32-bit) fields. Each record represents one depth and records are ordered by depth for each station. The 1st field of each record contains station code, the 2nd field contains depth. The remainder of the record is made up of six groups of 5 fields each which contain the five coefficients, A1, B1, A2, B2 and C respectively. The six groups represent the six data parameters temperature, salinity, oxygen, sigma-t, thermosteric anomaly and geopotential anomaly. A control card must be submitted at run time designating whether the output file is to be printed. The format is: (A6), and the entry "bPRINT" will invoke the print option. The deck set-up is shown in the following example.

# Example:

? BEGIN JOB
? RUN FIT/HARMONICS
? FILE OCNSEQ (TITLE=OCNSEQ/ALLCRUISES/LINE90)
? FILE HMCOEF (TITLE=HMCOEF/ALLCRUISES/LINE90)
? DATA
 (print control card)
bPRINT
? END JOB.

#### 9. HARMDEVS

This program computes the deviations of data values in OCNDAT or OCNSEQ (sorted by station code) files from corresponding values determined with harmonic coefficients computed by FIT/HARMONICS. All stations in the input file of data values must be represented in the file of harmonic coefficients. HMCOEF. The program identifies the first station on the input data file, then spaces through the reference file of coefficients until that station is found. The coefficients are used to compute a value for each data parameter at each depth and deviations from the corresponding data values are obtained. The deviations are either stored directly, as anomalies, on the output file, or accumulated for computing standard deviations, depending upon output options set at run time. If HARMDEVS is used to get anomalies, the input should be an OCNDAT file containing data from one or more cruises, each represented by a separate data set in the file. If it is used for standard deviations, the input should be an OCNSEQ file created by SORTOCNDAT with the data from all cruises sorted by station code in a single data set. The format of the output file, OCNDEV, will correspond to that of the input data file. A third output

option is available which stores the counts of data values accumulated for standard deviation in a second output file, OCNCNT. The format of the control card is: (2A6), where the 1st field is for a print option and the 2nd field is for the output option. If "bPRINT" is entered in the 1st field, the output file(s) will be printed. If the entry in the 2nd field is "bbANOM", the anomaly option will be invoked. If it is "bbSDEV", the standard deviation option will be invoked. If it is "bbCNTS" then both standard deviations and counts will be output. Parameter values computed with the harmonic coefficients will represent the date designated by the year-month-day codes in the corresponding data records. The deck set-up is shown in the following example.

# Example:

0.8N 37.3W 5207 5406 5508 5606 5807 5910 6001 6001 6002 21.6w 21.6w 5500 5807 5807 5907 5901 6004 6902 7205 20.8N 5.7W 5009 5107 5207 5508 5606 5807 126 50003 50004 50005 50003 51007 51009 5003 5004 5005 5005 5107 5108 5003 5005 5005 5006 5207 5207 5807 5910 6001 6902 7205 6004 6007 6902 7205 30.8N 30.5W 5107 5508 5606 5910 6001 20.8N 52.8W 124 50003 50005 50005 50007 50008 51007 51008 124 5003 5004 5005 5006 5008 CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978 46.8N 54.6W 55.1N 6001 6004 5.1N 55.9W 5107 5606 5804 5807 6004 5207 5606 5807 6004 6902 39 123 5108 5206 5206 125 124 124 5003 5005 5009 5009 5009 5004 5005 5008 5008 5107 15.1N 33.3W 30.4<u>N</u> 45.0<u>W</u> 50.4N 59.5W 5606 5807 5910 6004 6902 5606 5807 5910 6004 6902 7207 6902 7207 40 124 5207 5606 5804 5005 5007 5008 5009 5108 5005 5005 5007 5009 5107 44.7N 12.4W 5009 5107 6001 6902 24.7N 57.1W 5009 5107 5508 4.7N 27.4W 5806 5807 6001 6004 6902 7205 40 128 39 129 129 5000 5000 5000 5000 5000 5000 5004 5005 5006 5006 5005 5005 5006 4.7N 11.1W 5207 5606 5807 6001 6004 6902 24.7N 42.2W 44.7N 56.8W 5207 5606 5807 6001 6902 7205 5606 5807 6001 6902 7205 127 40 126 5004 5005 5008 5009 5107 5005 5005 5007 5008 5107 5005 5006 5006 5009 5107 TABLE 1. 5207 5606 5807 6004 6902 44.7M 39.1W 34.7N 2.2W 6902 7205 5606 5807 6004 24.7N 25.2W 5207 5606 5807 5005 5007 5008 5009 125 5004 5005 5007 5008 5107

CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

CON'T	6004	6101	6107	6110	6203	6210	6304	6401	6407 6501	6507	6601	6612	6801	6901	6902	6905	6910	6912	7203	7207	7503	7801	7805	7808							
08009	5008 5009	5105	5107	5108	5110	5111	5204	5206	5207 5209	5210	5211	5305	5307	5708 5405	5406	5407	5506	5510	5604 5606	5804	5807	5810 5901	5904	5910	1000						
s.		·· •- •			<del></del> .		-· <b>-</b> -·		<b></b>			· <b>-</b>		<b>-</b> -							<b>-</b> •			·· •-· •	- <b></b> -	,-		<b></b> .		<b>-</b>	
CON'T	6101	6110	6203	6210	6304	6401	6501	6507	6601 6607	6612	6801 6806	6901	6902	6907	6908	6910	7202	7203	7205	7412	7503	7803	7805	7808			79		3.2W		5006 5006 5007
02009	5104	5107	5109	5110	5112	5204	5207 5207	5209	5210 5211	5304	5305	5308	5506	2208 5510	5604	5606	5804	5807	5810	5904	5907	5910 6001	6004	6009			9009		125		5003 5004
CON'T	6201	6210	6304	6401	6501	6507	1099	6612	6801 6806	6901	6902 6905	1 2069	6908	6912	7202	7203	7207	7412	7501	7803	7804	7805	7808	•••••		ξ.	3	16.8N		5006	5008
09009	5207 5209	5240	5305	5306	5308	5404	5405 5406	5407	5408 5506	5510	5604 5604	5707	5801	5807	5810	5901	5905	5907	5910	6004	6007	6009 6101	6104	ò		02009	2,000	37	124	5002	5007 5004 5005
CON'T	6304	6407	6507	6601	6612	6801	6901	6902	2069 6907	6910	6912	7202	7203	7412	7801	7803	7805	7807	, 1808 1		•••		68	36.8N	ċ	5105	5107	5108	5110	5111	5204 5206
60052	5304 5305	5307	5506	5508	5707	5801	5804 5805	5806	5807 5810	5901	5907 5910	6001	6004	6009	6101	6104	6110	6201	6203	100			09009	37	C71	5002	5004	5005	500 <del>6</del>	5008	5101 5104
2	29.1N 21.9W	5910	6004	<b>-</b> , -		•	0		21.1N :	١ :	5807	6001	6004	2069		g •	9		11.1N		6004	7207			69	α	· ·		5207	5208	5210 5211
57051	38 123	5108	5207	5807			57055		38 123	1	5108 5206	5207	5406	2000			57060	1	38 124	-  -	5807	5910 6001			60052	7.4	123		5108	5110	5112
σ	0.8N 0.1W	5006	5008	2009	·· •-·	a-• •	t~		2.5N 52.0W	: :	5910	6004	• •			<b>-</b>	n	56.5N	•	5910	6001	6904				9	•	27.4W	6004	6902	0 0 0
50130	130	5002	5004	5005			53052		39 123		5108 5206	5406	5807			22052	CCOCC	38	124	5108	5206	5207 5606	5807			53060	38	124	5807	5910	000
CONT	6902				2	0	•	5008	5009	6001	6902 7205				6	- 0	33.3W		5007	5009	5107			··· <del>-</del> -· •	_	20.8N	. <u>.</u> .	5008	5508	6902	502)
50090	5910	9009		0	20.00	38	12.	5005	5003	5005	5006				50110	1	128		5002 5002 7003	5007	5005	2006		6	50120	37	2	5002	5007 5004	5005	5007

38
<u></u>
6
195
•
NG
URING
70
SE
Ë
CRUISE
MORE
Ξ
OR
: :
SIX.
NO
Θ
臣
Π
OCCUPIE
0
NS
10
ΑŢ
ST
H H
COF
길
$\ddot{\circ}$

63057 10	37 8.6N 122 58.7W	5006 5105 5007 5107 5008 5109	101 5	•	0 1	123 11.7W	707	807	901	907	910 001	004	101	107	! 6201 7412 ! 6203 7501	210 301	304	407	507	<b></b> (0.00)	<b></b>	<b></b>
63052 CON'T	7203 7803 7207 7804 7412 7805	7501 7807 7503 7808 7801		رح ا	37 12.6N 122 50.1W	9 80	110	112 6	205	207	208 209 605	210 6	304 6 305 6	306 307 307	308 6 404 6	405 7 406 7	510 7	200	6009 7801 6101 7803	104	6201 7808 1 6201 7808	6029
61055 15	37 36.5N 123 7.2W	002 003 51 004 51	5005 5006 5007 5007 5109	600		63052 67	37 18.6N	90	+ + 0	204	205 206	207 208	209	211	305 306	307 308	407	510	801	807	5901 6910 5904 6912	907
60160 14	34 16.8N 130 42.6W	07 62 09 62 01 63	6104 6304 6107 6407 6110 6501	5	<b></b>	9 2 2	132 5.8W	007 62	101 63	107 64				60200 13	32 56.8N 133 28.2W	69 70	00 62	6104 6304	רו רו			
60120 27	35 36.8N 127 54.3W	002 003 004 6	5005 6201 5006 6203 5007 6210	900	107 6 305 6	007 009 7	2		60130 9	9	8 36.	002	5004 5009 1 5005 5106 1	5006		50140	)	129 18.8W	6007 62	6101 63	6110 6501	6201 65
60100 45	36 16.8N 126 29.1W	02 54 03 58 04 58	5005 6001 5006 6007 5007 6009	09 04 61	05 61 06 62	07 62 11 63	06 64	10 65	05 66	69 80	27 70	-		60110 17	35 56.8N 1	002 5105	003 5106	005 5209 005 5209	5007 5211 1 5008 5305 1	009 5804	-	
£ 42 06009	36 36.8N :	002 59 003 59 004 59	5005 5006 6004 5007 6007	009 009 104 61	105 61 106 61	107 62	111 63	204 207 207 207 207	206 65	209 662	210 66	304 68	307 69 308 69	506 69	604 606 606 606	804 72 805 72	906 75	810 78	904			pe.

5904 5907 5910 6004 6908 6912 7203 7203 7801 7805 7805 7805 7805 51112 52112 5203 5209 5209 5304 5407 5408 5510 5803 5804 5901 5904 5904 5907 5910 6907 6905 6905 6908 6912 7202 7203 7204 7412 7501 7801 7803 7803 7805 7805 47.2N 11.7W 6908 6912 7203 7203 7412 7803 7803 7803 7805 77.20 78.00 7207 7412 7412 7503 7510 7803 7804 7805 7804 7805 7805 7.2N 7.8W 6801 6800 6901 6902 6907 6907 6910 7201 7202 5000 66501 66501 66501 66501 66901 6 6107 6201 6201 6201 6304 6401 6407 6601 5804 5804 5807 5904 5907 5910 6004 6009 7.0000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.000 7.00000 7.0000 7.0000 7.0000 7.0000 7.0000 7.0000 7.0000 7. 66501 66501 66601 66601 66901 66902 77202 77203 7 5205 5206 5206 5209 5210 5210 5211 5205 5206 5207 5207 725075 72 6903 6910 6912 7203 7203 7203 7801 7805 2.6N 20.5W 6907 6908 6912 7203 7203 7503 7805 7805 5204 5205 5206 5807 5807 5810 5905 5907 5910 6001 6607 6801 6801 125 5904 5904 5907 6004 6801 6801 6901 6901 5109 5110 5111 124 48.6N 41.9W 5604 5606 6912 7202 5106 5107 42.6N 54.8W 52.6N 33.3W 123 5807 5807 5901 6004 6602 6801 6902 6902 5205 5206 5207 5305 5007 5101 5104

CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

CALCOFI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

70110 19 34 12.9N 125 52.6W 5002 5105 5004 5107 5005 5508 5007 5804 5008 6902 5009 6905 5104	70120 25 126 34.2W 5003 6104 5004 6107 5006 6201 5006 6201 5007 6203 5009 6210 5009 6301 5009 6301 5106 6401 5107 6501 6101 533 32.9W 127 15.6W 5004 5008 5006 5010
6704 6908 6401 6910 6401 6910 6407 6912 6501 7201 6601 7202 6601 7203 6601 7203 6601 7203 6801 7803 6901 7803 6902 7805 6905 7807	70100 43 124 32.9N 125 10.8W 5002 6007 5003 6009 5004 6101 5008 6201 5009 6201 5104 6301 5104 6301 5107 6407 5205 6601 5305 6601 5305 6601 5305 6601 5305 6601 5305 6601 5305 6601 5305 6601
70080 CON 'T 6501 7202 6501 7203 6601 7203 6601 7207 6801 7503 6806 7507 6901 7507 6902 7801 6908 7805 6908 7805 6910 7801 6912 7801	70090 78 78 78 78 78 78 78 78 78 78 78 78 78
ρ -4	5002 5004 5005 5004 5005 5006 5006 5006 5006 5009 5009 5104 5104 5105 5107 5108 5107 5108 5107 5108 5109
0 ",	5008 5904 5009 5907 5010 5910 5104 6004 5105 6007 5106 6009 5107 6101 5110 6110 5111 6201 5204 6210 5204 6210 5208 6407 5208 6407 5208 6407 5208 6605 5305 6801 5305 6801 5508 6905 5707 6910
25 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	5107 5108 5108 5109 5110 5111 5111 5204 6203 5204 6501 5204 6501 5209 6601 5209 6601 5209 6601 5209 6801 5209 6801 5304 6902 5304 6901 5406 5707 6908 5707 6908 5707 5707 6908 5707 5707 6908 5707 5707 5707 5707 5708 6912 5707
25.2 25.2 25.2 25.2 25.2 25.2 25.2 25.2	5204 5205 5205 5206 5306 6801 5305 6801 5305 6901 5504 5508 5508 5508 5707 7202 5707 7202 5606 7202 5606 7203 6001 7803 6004 7803 6009 7804 6009 7805 6009 7807 7807 7807 6009 7807 6009 7807 7807 7808 6009 7808 6009 7809 6009 7809 6009 7809 6009 7809 6009 7809 6009 7809 7809 6009 7809 6009 7809 6009 7809 6009 7809 7809 7809 6009 7809 7809 7809 6009 7809 7809 6009 7809 7809 7809 7809 7809 6009 7809 7809 6009 7809 7809 6009 7809

5105 5106 5107 57.3N 3.5W 66801 66801 66901 66902 66903 77202 77202 77202 77203 7 6002 6401 CON 'T 5007 5008 5101 5902 5911 5912 6908 6910 6912 7201 7202 7203 7203 7801 7801 7807 7202 7203 7412 7801 7803 7805 7805 5109 5110 52112 5204 5205 5206 66004 6901 6905 6907 6908 6910 38.6N 21.9W 5907 5910 6001 6902 6905 6907 6908 6910 7201 7203 7412 7412 7804 7804 7805 7805 16.6N 1.9W 5106 5107 122 5804 5807 5810 5007 5101 5104 CALCOFI STATIONS OCCUPIED ON 61010 61010 61010 61010 62011 6 CON 'T 6001 6002 38.6N 5300 5300 5300 5300 5300 7202 7202 7202 36.6N 5105 5107 5711 6203 6210 6301 6304 6507 2.9N 70130 CON'T 5007 5008 5101 51112 52112 5204 5205 5206 5208 5208 6009 6101 6104 6107 

SIX OR MORE CRUISES DURING 1950-1978

CON 'T

m
6
<u>ص</u>
Ţ
8
9
•
9
E
黑
ಗ
m
βĄ
ESI
Þ
33
贸
8
Σ
3
_
SIX
NC
0
A
E
딢
CUP
ರ
0
ß
SNO
Н
¥
턵
Ø
띥
0
S
٧
O

CON'T	5904 5905 5905	5907 5908	5910 5911	5912 6001	6002	6003 6003 7	9009	6007	6101	6102	6107	6110	6203	6207	6210	6304	6307	6210	6407	6501	6601	2099	6612	6801	9089	6901	2069	6904 4004	6907	6908	6910	7201	7202
80060	5109	5112	5202 5203	5204 5205	5206	520.7 5208	5209	5210	5301	5302	5304	5305 5305	5307	5308	5502 5504	5506	5508	77.0	5602	5604	5704	5707	5710	5712	5801	5802	7807	7804 7805	5806	5807	5810 1810	5901	5902
		· <b>-</b> · - ·					· <b>-</b>				. 🛶	<b>-</b> · -	<b></b> .		<b></b>	<b></b> .	<b></b>		••••								-· <del>-</del> -	•-	. <b>.</b>	<b>-</b> ·	<b></b>	. <b></b> .	<b>-</b>
CON'T	6304 6307 6310	6401 6407	6501 6507	660 <b>1</b> 6607	6612	6707	9089	6901 6902	6904	6905	6908	6910	7201	7202	7203	7501	7801	7807	7805	7807	200			123		0.00 NO.00	•				5104		
80055	5801 5802 5803	5804 5805	5806 5807	5810 5811	5812	7901 7901	5904	5907 5908	5910	5911 5910	6001	6002	6007	6009	6101	6104	6107	6201	6203	6210				80060		34	121	5002	5003	5004	5005 5006	5007	5008
			<b></b>				. <b></b> .				. <del>-</del>		<b></b> .			· <b>-</b>		<b>-</b> -	<b></b>					· •	<b>-</b>	<b>-</b> . •	<b>-</b> -	<u>.</u> .	. <b>.</b>	<b>-</b>		· <b>-</b> -	
-	23.0N 39.8W	6207 6401	6407 7202	7412				124	19.0N	œ	5302	5303	5305	5306	5307	5401	5402	5405 5404	5405	5406	5410	5502	5504	5508	5510	5511	2144 2007	7007	5606	5704	5707	5711	5712
80053	34 120	5902 5911	5912 6001	6002	)			80055	34	120	5005	5003	5005	2006	5007 5008	5101	5102	7017 7007	5105	5106	5109	5110	5111	7.50 1000	5202	5203	5204 5205	7000 7000	5207	5208	5209	5211	5301
					<b></b> -							<b></b>	<b></b> .			. <b>.</b>			<b></b>		<b>.</b>	<b>-</b> ·	<b>-</b> - •		<b>.</b>	<b>-</b>		-· <b>-</b> -	<b></b> .				<b>-</b>
8	25.0N 35.6W	6009	6102 6104	6107	6201	6203 6210	6301	6304	6310	6401	6501	6507	6607	6612	6707	6806	6901	2069	6904	6905	2069 2069	6910	6912	7201	7202	7202	7203	7410	7412	7801	7803	7807	7808
80052	120	5108 5109	5110	5112	5202	5203	5204	5205 5205	5207	5208 5208	5210	5211	5302	5303	530 <b>4</b> 5305	5306	5307	7.70g	5506	5510	5712	5801	5803	7804 5805	5806	5807	5810	701 701	5901	5904	5907 5908	5910	6007
							. <b>-</b> .		. <b>-</b> .	<b>-</b> · -	. <b>.</b> .		<b>.</b>			٠ ـ.	<b>-</b>	-· <b>-</b> -	. <b>-</b> .		- · <del>-</del> ·		<b>-</b> ·•			•		<del>-</del>	· <b>-</b>				
31	3.3N 56.5W	6907 6908	6910 6912	7201	7203	7412	7510	7801 7803	7805	7807				1	<b>58</b>	43.3N	ά	7009	6905	6907	6910 6910	6912	7201	7007	7801	7803	7805	78087					
77080	34 122	5506 5707	5804 5807	200	8	£8	88	96	88	86	88			i	77090	33		5506	5707	5803	5804 5807	5810	5904	7000 7010	6004	6601	6607	0690	0 0 0 0				
			<b>-</b> ··-·		• ••••		<b></b> .		· <del>-</del> -	<b>-</b>	· <b>-</b> ·		<b>.</b>				<b></b> •	~· •-	• •	<b></b> -	<b>-</b>					<b>-</b> ···		<b>-</b> -	· <b>-</b> -			· <b>-</b>	
30	33.3N 53.9W	$\sim$	5210 5211	N K	/ <b>K</b> U	Мκ	ľ	ဖင	7	$\alpha \circ$	J			37	K	14.8W	3	- G	6902	88	38	5	200		20	50	80	2 6	88	80	8		
77065	34	5006	5101	5105	5107	5108 5109	5110	5111	5205	5206	760			77070		122		20	5804	ထင	ρσ	$^{\prime}$	$\sigma$	νσ	ıΦ	$\sigma$	$\sigma$ c	$\nu$ C	$\circ$	4	44	2	
							··																						. <b>-</b> .			· <del>-</del> ·	
16	49.3N 20.3W	6110	6203 6210	6301	6501	6507			47	K	32.94	2402	6601	6607	6612	6806	6901	690v	6907	3069	6912	7201	7202	7007	7801	7807	7804	7007	78087				
77057	121	5203 5204	5902 6001	6002	6104	6107			77060		12,	¥000	5204	5707	5801 5801	5804	5805	7807 7807	5901	5902	5904	5908	5910	100 100 100	6001	6002	6003	9004 9004	6010	6401			

5412 5604 5606 5704 5707 5804 5807 5807 5807 5807 5807 5103 5104 5304 5305 5804 6207 29.0N 37.2W 6210 6301 6304 6407 6501 7300 7300 7300 7404 7404 7404 7408 7408 5003 5004 5005 5006 5006 6101 6104 6107 6110 6201 6104 6107 6110 6201 6203 6203 6204 6201 6301 6304 5201 5201 5304 5305 5305 5305 5305 5305 9.0N 6.1W CON'T 125 49.0N 54.4W 7201 7202 7203 7205 7207 7201 7201 7801 7803 7803 7808 CON . T 5000 6601 6607 6612 6707 6806 6901 6902 6905 6907 6908 6908  $\begin{array}{c} 0.000 \\$ CON'T 29.0N 32.0W CON'T 59807 59807 59807 59807 59807 69 7803 7804 7805 7807 CONT 49.0N 50.6W 5005 7205 7207 7412 7501 7507 

SIX OR MORE CRUISES DURING 1950-1978

Ö

CALCOFI STATIONS OCCUPIED

CALCOFI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

TO CON'T	808		80 62	32 54.7N		002 5907	7,17	·ω	w.	t) «	Œ	Θ	w (	D C	, w	6	6	o w	Φ	ω (	0 5	- [-	. [~	<u></u> 1	[	- [-	. [-		[_						
83070	78		83080	•	-	50	7 2 2 2 2	50	50	7 2 2 2 2	3.5	5	(C)	ン R ソ ひ	150	33	10 I	2 4 4 4	75	R R	シ で ひん	500	57	57	ر ا	טית מַמַ	200	59	J 9	r. Q					
							-· <b>-</b>	· <b>-</b> .		<b>-</b> -	. <b>-</b>		<b>.</b>		<b></b>			<del>-</del> -	· <b>-</b> ·	•	-· <b>-</b> -	·	<b>-</b> ·	<b>-</b> ···	<b></b> •	-· <b>-</b> -	. <b>.</b> .		<b>-</b> .	<b>-</b>	· <b>-</b> -				<b>-</b>
CON	7510	7803	7805	7808			83		14.7N	ċ	5910	6001	6004	6501 6507	6601	2099	6612	6801	9089	6901	2000	6905	6907	6908	6910	7207	7202	7203	7412	7501	7712	7801	7803 7804	7805	7807
83060	6905 6907 6908	6910 6912	7201	7203			83070		K. 60		5005	5004	5005	7006 7007	5008	5011	5101	5107	5204	5205	7206	5305	5306	5504	5506	7604	5704	5707	5710	5801 5807	5810	5901	5902 5904	5907	5908
	•			· · •	-· <b>-</b> -		-· <b>-</b>	· <b>-</b> -			. <b>.</b> .				<b></b> .			·· •·		<b> •</b>	-· <b>-</b> -	· <del>-</del> ·		<b>-</b> ···	•						·			• •	
106	34.7N 45.3W	5801 5804																																	
83060	33 120	5002 5004	5005	5007	5011	5101	5102 5104	5105	5110	2204 2004	5206	5301	5302	2007 2008 1404	5305	9266	5307	5311	5312	5401	7407 7407	5404	5405	5406	7408	7410	5502	5506	5500	7510 7510	5512	5602	7604 7606	5704	5707
<b></b> -					·· •·		-· <b>-</b> -	. 🛶	<b>-</b> . •		<b></b> .	⊷.		<b>-</b> -	· <b>-</b>	<b>-</b>		<b>.</b>	<b>.</b> .	<b>-</b>	-· <b>-</b> -			<b></b> ··		<del>-</del> -	. <b>-</b> .		<b>-</b>	<b>-</b> ·-	. 🕳				
CONT	7807 7808		ለ	-	24.6W		7907 7907	5904	5907	7908 7910	6001	6004	6401	6501	6601	2099	6612	6801	9089	6901	20.00 20.00 20.00	6908	6912	7201	2027	7507	7712	7801	7803	7804	7808				
83051	7803 7804		83055	) <b>!</b>	120	0	700r	5005	5006	7007	5102	5104	5105	71.00 0.00	5110	5204	5205 5205	5301	5302	5303	7304	5306	5307	5308	5509	5311	5312	5604	5606	5801 5801	5807	5810			
				. <b>.</b>				. <b>.</b> .	<b>-</b> · .		<b></b> .			<b>-</b> -	<b></b> .	<b>-</b> ·		·· •	<b>-</b> .		<b>-</b> -	· <b>-</b> ·			<b>-</b>	·· <b>-</b> -	<b></b>								
12	58.7N 55.5W	5307 5308	MM	MM	Ω.			64	(	30.0 0.0 0.0	•	5810	5901	700c	5907	5908	5910	6004	6401	6501	6601	6607	6612	6707	6801	6901	6902	6904	6905	6907 6908	6910	6912	7202	7712	7801
83048	119	5301	K K	/ IO/ N	Α.			83051	1	120	) 	5011	5301	7,002	5304	5305	5306	5308	5309	5310	- 100 - 100 - 0180	5401	5402	5403	7404 7.08	5400	5407	5408	5410	5412 5502	5504	5510	5512	5804	5807
				·	· •-·			· <b>-</b> ·	<b></b>	-• <del>-</del> -					<b>-</b> .							. <b></b> .	<u>.</u> .		<u>-</u>			<b>-</b>			<b></b>	<b>-</b> ···	-· <b>-</b> ·		
75	8.7N 34.7W	5804 5807	5810 5811	5901	5904	5907	7970 7910	5911	5912	6002	6009	6004	6009	6501	6507	6601	6607	5100	6801	6806	0009	6904	6905	2069	2000	6912	7201	7202	7602						
83043	34	5111		K / K	1 14 /	$N \setminus N$	) K	/ PK /	$\sim 1$	) N)	· N \	N 1	-	+ •		*+ *	-	+ <+	44	<b>*</b> + 10	$\neg$	$\sim$	$-10^{\circ}$	$\sim$	$\cap \iota \iota$	310	_	~ 1	_	m					
										·· •-·			-· -			<b>-</b> ··•		. 🛶							-· <b>-</b> -	· -		<b>-</b> ···	<b>-</b>		<b></b>	<b>-</b> ···		<b>.</b>	
CONT	6801 6806 6901		w w	w u	<b>.</b>	(~ t	- [-	- [	[- t	- [-	[-					-	_	22.2W		5502	νц.	۱u ۱	(A					80	,	10.7N 30.5W	•	7804	7807	7808	
82047	5904 5907 5908	5910 5911	6001 6002	6003	6005	9009	6407	6501	6507	6607	6612	6707				83040	7.4	119		5302	3,0	30	30	30				83042		119	`	4	7602	$\infty$	

9.1M 9.1W 9.

87040 CON'T \$\text{\tt}\xint{\text{\ 14086  $\frac{660}{6}$ OR MORE CRUISES DURING 1950-1978 55215 55801 55801 55801 55801 55801 569 7801 7803 7804 7807 7808 0000 118 5604 5604 5704 5707 5402 5404 5405 5405 5406 5412 6412 57.0N 14.3W 5203 5207 5302 5303 5305 5305 5309 5311 5311 5510 55110 5520 5520 5520 5520 55210 55210 121 5108 5201 SIX CALCOFI STATIONS OCCUPIED ON 5403 5403 5403 5406 5406 5410 5410 37.0N 51.8W 5302 5303 5304 5305 5306 5307 5311 5311 5311 47.0N 31.1W 5300 5300 5300 5300 5300 5311 5311 5404 5404 5404 5408 5410 5511 55203 5301 5301 5300 5300 5300 5310 5311 7602 5302 5303 5303 5303 5303 5312 5312 5402 119 66001 6 5207 5208 5209 5210 5211 5110 5110 5112 5201 5202 

2002 1 73 2 1 1 73 2 1 2 75 2 2 2 100 2 200 2 2 00 2 2 0 0 2 2 00 2 2 0 0 2 2 0 0 2 2 0 0 2 2 1 0 0 0 2 2 0 0 2 2 1 0 0 0 2 2 0 0 2 2 1 0 0 0 2 2 0 0 2 2 1 0 0 0 2 2 0 0 2 2 1 0 0 0 2 2 0 0 2 2 1 0 0 0 2 2 0 0 2 2 1 0 0 0 0 2 2 0 0 2 2 1 0 0 0 0 2 2 0 0 2 2 1 0 0 0 0 2 2 0 0 2 2 1 0 0 0 0 2 2 0 0 2 2 1 0 0 0 0 2 2 0 0 2 2 1 0 0 0 0 2 2 0 0 2 2 1 0 0 0 0 2 2 0 0 2 2 1 0 0 0 0 2 2 0 0 2 2 1 0 0 0 0 0 0 0 0 0 2 2 1 0 0 0 0 0 0 0 0 2 2 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	).
90028 CON'T 5708 6005 5710 6005 6005 5710 6005 6005 5710 6005 6005 5710 6005 6005 5710 6005 6005 5710 6005 6005 5710	
97090 51 22 25.4M 52002 6004 50004 6004 50005 6507 50005 6507 5101 6801 5104 6904 5204 6908 5204 6908 5204 6908 5204 6908 5205 6908 5206 6908 5207 7707 5208 6908 5208 7807	202
7201 7801 7202 7804 7412 7803 7712 7804 7712 7804 7712 7805 7712 7805 7712 7805 7704 7807 7704 7203 7704 6501 7704 6501 7704 7701 7704 7701 7707 7701	
87060 CON-T 6004 6908 6005 6910 6504 7201 6504 7202 6607 7202 6801 7412 6801 7712 6904 7801 6904 7804 6904 7804 7703 6904 7804 7703 7703 7703 7703 7703 7703 7703 7703 7703 7703 7704 7704 7703 7703 7703 7704 7704 7704 7704 7704 7705 7705 7705 7706 7706 7707	801 6
87055 CON.T. 5801 5902 CON.T. 5804 5900 CON.T. 5804 5900 CON.T. 5902 7201 6801 CON.T. 5902 7201 6801 CON.T. 5902 7201 6801 CON.T. 5902 7202 CON.T. 5904 5901 CON.T. 5906 5901 CON.T.	410 6
87050 119 39.4N 119 39.4N 50002 55002 5101 39.4N 52002 55004 5101 5510 52004 6501 52004 6501 53004 6501 53004 6501 53004 6501 53004 6501 53006 69004 53007 5506 53007 6900 53007 5500 54007 7202 54007 7203 55007 7203	306 570

CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

CON'T	7210	7801	7804	7802	7808	•			126	<u>.</u>	57.6W		5301	5502	7207 7304	5305	5306	5307	7208	5312	5502	5504	5506	5510	2177 2003	5604	9099	5704	5707	5710	7 - 0	7807 7803	5807	5809	5810	781-	5902
90053	6905 6907 6908	325	201	200	88				09006	0	119		5002	5004	7000r	5007	5008	5006	1010	5103	5104	5105	5106	5107	7 10 10 10	5110	5111	5112	5201	5202	7070 7000	7007 7007	5206	5207	5208	7707 7 <b>21</b> 0	5211
					<b>-</b>			·· -·			<b></b>	<b>-</b> .		•		. <b>.</b> .	<b>-</b> .	<b>-</b> ··	-•	<b>-</b>		<b>-</b>		<b>-</b> ··•		·				-· -	-· •	<del>-</del> -	· <del>-</del> ·	<b>-</b> .	<b>-</b> - •	<b>-</b> -	• •
CON'T	5812 5901	5903	5904 5905	5906	7907 7908	5910	5.91 1.01	6001	6002	0000	6005	9009	6007	8009	200	6104	6107	6110	6201	6211 6211	6302	6304	6307	6310	5401	6407	6410	6501	6504	6507	7099	6612	6707	6801	9089	- 000 - 000 - 000	6904
90053	5111	5202	5204	5205	5207	5208	5209	5211	5301	5502 5202	5304	5205	5306	5507	7700 7710	5312	5401	5402	2405	2404 5405	5406	5407	5408	5506	7508 7508	5510	5511	5512	5602	5604	2000	5710	5712	5802	5803	780.7 7810.	5811
CON'T	7805				cc	45.1N	9	5901	5902	7903 2004	5907	5908	5909	5910	1010	6001	6002	6009	6004 	2009	6207	<del>-</del> ,		•		132		39.1N	œ	-			-	_	5107		
90045	7801 7803 7804	2		i G	90050	32	119	5303	5304	7,507 7,007	5307	5308	5310	25.72	0.4.C	5502	5504	5712	7807	5809 5810	5811	5812				90053		32	6	C	2002	5007	5005	5006	5007	200g	5101
CONTE	6002	6005	2009	8009	6101	6104	6105	6110	6201	6203	6302	6304	6307	6510	1040	6407	6410	6501	6504	6607	6610	6612	1 2029	6801	6806	6902	6904	6905	1 2009	6908	07.0	7202	7203	7205	7207	7602	7712
90045	5210 5211	5302	5304	5305	5307	5308	5310	5502	5504	7706 7106	5512	5605	5604	2606 7400	5704 5704	5707	5710	5712	2802	5807 5804	5805	5806	5807	5809	7810	5812	5901	5902	5903	5904 1005	700r	7909 5907	5908	5909	5910	- 120 - 0	6001
CONT	7207	7602	7801	7803	7805	7807	1808				7		7.1N	٠	S	6910	6912	<b>6</b>		•• •••		129		55.1N	ċ	5108	5109	5110	5111	5112	1020	2003	5204	5205	5206	7000	5209
90037	6806 6901 6903	88	383	88	2.2	20	200	2			90039				ç	8	6905	9				90045		32		00	00	ò	8	ŠŠ	38		9	0	5103	ŠČ	0
<u>.</u>				•				<b>-</b> .			• •				·· <del>-</del>				-· <b>-</b>					<b></b>					•			··					• =-
CON'T	5809 5810 5811	5812	5905	7903	790 900	5906	5907	5900	5910	200	600	6002	6003	9000	000	6007	6008	6010		6107	6110	6201	6203	6211	2000	6307	6310	6401	6404	6407	04-0	6504	6507	6607	6610	6707	6801
90037	5201 5202 5203	5204	5206	520.7	5209	5211	5301	5303	5304	7207 7306	5307	5308	5310	2166	5407	5403	5404	5405	2400	5407	5410	5412	5502	5504	7700 7500	5510	5512	5602	5604	2606 700	7070	5707	5710	5712	5802	5804 5804	5807
								· -·			· <b>-</b> -		•		·· <u>-</u>	٠ ــ.			-· -	<b>.</b>		<b>-</b>				· <u>-</u> .	<b>-</b>		•		<b>-</b> -	· -			<b>.</b>		<b></b>
CON'T	6801 6806 6904	888	283	38	32	5	200	20	200	N				t	_	•	₩.9	i C	1807	7808					130			ķ.	(	$\frac{1}{2}$	2 5	20	5	5	5109	- =	=
90032	6211 6302 6304	/W/N	U 44 .	♥ ・	ਰ ਦ	. CO 1	ഥവ	വ	SO.	Ω Γ	-			22000	3000		118	i	777	7803	7804				75000		33		•	$\circ$	) C	$\circ$	$\circ$	0	5008	> ←	-

28
φ.
7
Ċ
6
•
Ö
E
DUR IN
-
E
50
CRUISE
R
O
Œ
$\Xi$
MORE
SIX OR
X
H 70
U.
8
NO 0
$\boldsymbol{\omega}$
Ξ
d fi
5
OCCUPIE
ONS
0
Ē
STATI
STATIONS
Œ
COPI
-
A C
_

28	45.1N 19.9W	5107	5305	5401	7900 000	6207	6507	6610	6612	10/0	7807	7808				45		25.1N	,	6110	6201	6203	6207	0720	6304	6401	6407	6501 6507	- 679	6612	6707	6901	6904	2000	1000	7210	7804	7807
90110	30	88	5004	88	88	38	8	5	0;		50	0				90120		30	(7)	5002	5003	5004	5005	5000	5008	5009	5101	2017 2018	200	5105	5304	5305	5508	7909	2000	6101	6104	6107
			<b>-</b> .	<b>-</b>				<b>-</b> .			. <b>.</b> .	<b>-</b>	<b>-</b>	-· •	<b>-</b> -	. <b>.</b> .			<u>-</u>		<b>-</b>					<b>-</b>	<b>-</b> ·						<b>-</b> ··	-· -	<b>-</b>	<u>.</u>	<b></b>	
29	5.1N	6104	6110	6201	6203	6010	6301	6304	6401	6404	6410	6501	6504	2000	6607	6707	6801	6901	4000	6910	7205	7210	7412	7007 1007	7803	7804	7807	7808										
90100	31 122 3	200	5007	005	006	٠ a	600	101	102	507	105	106	107	200	204 206	207	304	305	900	910	001	005	003	004 005	900	200	800	010										
																			<b>-</b>		. 🕳 .			-· <b>-</b>	<b></b> .	<b>.</b>						<b>.</b>		<b>.</b>				
CONT	6104 6107	5201	5207	5210	5301	5204	5310	6401	5404	5407	5501	6504	5507	2.099	6610	2102	6801	9089	1069	2009 2004	6905	2069	6908	6910	7007	7203	7205	7207	0.50	7501	7801	7803	7804	7805	7807	808/		
06006	5205 5206	2010	77.7	53(	53(	у 4 6	54.0	55(	55	יל הליל	76 76(	57(	57	i	ر بر م	200	58	28	ž į	ນິດ	50	50	50	n o	n n v o	50,6	59	Š	000	ŠČ	Š 9	Ğ	Ğ	Ğ,	3	61		
<b></b>				<b></b> .	<b></b> .		· · -	. <b>.</b>		<b></b> •	<del>-</del> -			<b>.</b>		. <b>-</b>	<b>.</b> .		•	<b>-</b> -	. <b>.</b>	<b>-</b> ·			<b>-</b> -		<u>.</u> .	<b></b>	<b>.</b>					<b>-</b>	•	-· <b>-</b> -	. <b>-</b> .	
E, NO	6610 6612	6707 6801 6801	6901	6902	6904	6907	6908	6910	6912	7202	7205	7207	7210	7412	7501	7510	7712	7801	7803	7804 7805	7807	7808				110		5.1		5103	5104	5105	5106	5107	5108	5201 5202	5203	5204
Ö	6001 6002																									0		21.2		Š	1 60	40	05	90	0,4	800	95	02
90080	909	99	99	9	9	2 2 7	<u>.</u> .	5	9	62	200	62	63	63	60,4	9	9	64	9 6	ט ע נית	920	99				06006	,	•	_	נ	) C	50,0	50	20	ည်	200	3.22	7.
					<b>-</b>		<b>-</b>	. <b>.</b>	<b>-</b> ·	<b>-</b> ··	-· <b>-</b> -			<b></b>	<b>-</b>	=-				-· <del>-</del> -				<b>-</b>	-· <b>-</b> -			<b>-</b> . •				. <b>-</b> -	<b>.</b>	<b>-</b>	<b>-</b>	<b></b> -		
T, NO	7205	7412	7712	7801	7803	7804	7807	7808				112		.5.1N	œ	5504	5510	5512	5602	5604 5606	5704	5707	5710	5712	5801 5803	5807	5810	5811	5812	5902 5 <b>90</b> 3	7000 4004	5905	5906	5907	5908	5909 1010	5911	5912
70 ت	801											80		31 4	21	S	40	5006	20	α 2 C	0.0	02	60	0 1	200	04	08	50	210	200	ر د د	90	201	<u>0</u>	205	203	505	306
9007	68 68	600	6,6	69	69	9	70	72				90080			-	r C	200	50	ည်	ν. Ο C	7.5	12	5	ני ני	ر د	7.00	7	52	30	ν. Ω 0	. rc	, <b>r</b> C	52	FC	5	ריט ה ה∵ ה	ب ال	(TC)
<b></b> .							<b>-</b> -	. <b>-</b> .		<b>-</b>		<b></b> .	<b>-</b> ·		<b>-·</b> -	<b>-</b> -	· -					<b>.</b> .	<b>-</b> ·					<b>.</b>		-· <b>-</b>		. <b>.</b>		⊷.				
E.	50	60	- ~	25	50	4.0	ر ر د م	20	98	60	o <del>-</del>	2	5	02	03 03	و ر 4 بر	90	20	80	ے د	- 70	20	0	50	200	- -	02	04	7.0	25	5 8	07	10	5	04	07	- 6	0.7
CON	5803 5807																																					
90070	5106	5108 5109	5110	5201	5202	5203	7204 7205	5206	5207	5208	5209	5211	5301	5302	5303	7204 7305	5306	5307	5308	5510	5402	5403	5404	5405	2406 FA05	5408	5410	5412	5502	5506	20.00	5602	5604	2606	5704	5707	5712	5802
•••	<u></u>	<b></b>					<b>-</b> -					. <b>.</b>	٠.	<b>-</b> ·	-· •	<b>-</b> -			<b>-</b>			<b></b> .	<b>-</b> ·		-· <b>-</b>		<b>-</b> ·				·· -						·· •-	
CON 'T	6407 6410	6501 6504	6507	6610	6612	6707	6801 6804	6901	6902	6904	6905	6908	6910	6912	7202	7207 7205	7207	7210	7501	7267	7801	7803	7804	7805	1,087	000			1	132		38.3W		$\circ$	<b>-</b>	5102		-
09006	5903 5904	5905 5906	5907 5908	5909	5910	5911	2160 2003	6002	6003	6004	6005	6007	6008	6010	6101	6104	6110	6201	6203	6207	6302	6304	6307	6310	6401	4040				90070	32	120		5005	5004	5005	5007	5008

73008 73009 73 CON'T 7412 7412 7412 7712 7801 7803 7804 7805 7805 7805 7805 6901 6902 6904 6907 6908 6910 6910 7202 5105 5105 5105 5109 5109 6806 6901 6902 6907 6907 6910 6912 7202 5812 5902 5903 5905 5905 5910 5911 6002 6003 5810 6907 6909 6909 6910 6912 6912 7202 7202 7203 7808 7801 7803 7804 7807 7808 2107 2107 2107 2108 2108 6501 6604 6604 6607 6610 6610 6801 6806 6902 6902 6902 6002 6006 6102 6107 6110 5003 5003 5005 5005 5101 CON 'T 5812 5812 5902 5002 5002 5002 5002 6002 6002 6002 6002 6002 6003 6501 6507 6210 6301 6304 6407 6407 6501 6507 7804 7808 25.1N 57.3W 6401 6401 6501 6501 7804 7804 6203 6207 6210 6301 6304 5304 5305 6007 6010 6101 6107 6110 6203 127 6010 6101 6104 6107 6110 6201 6207 6010 6101 6104 6107 6707 7804 7807 7808 90120 CON'T 5.1N 59.8W 6507 6612 6707 6904 6904 6907 6910 7210 7210 7205 7210 124 5304 5305 5804 5905 6207 60100 61010 61004 61100 6201 6201 6201 6201 6401 6401

CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

CONT	6901 6902	6905	6908	6910	7202	7203	7501	7801	7803	7805	7807	909				39	Ċ	15.5W	,	6404	04-0 07-07-0	6604	6610	6612	6707	000	6904	2069	6910	7202	7412	7804	7804	7807	808/		
93090	6203	6302	6401	6404	6504	6507	6601 6604	6607	6610 6612	6707	6801	9099				93100	C	122	C C	7,000 0.000 0.000	2912 6002	6003	6004	6005	9009	7009	6010	6102	6104	6107	6110	1020	6207	6210	2059	1040	
		·				<b>-</b> ···		· ·	<b></b>		<b>-</b> ·•	-· <b>-</b> -		· <b>-</b>		<b></b>	<b>-</b>	<b>-</b> -	<b>-</b> ··	<b>-</b>	- · -	· <b>-</b> -				<b>-</b>	-		<b>-</b> ·	<b>-</b> ·	<b></b>	<b>.</b> .				·· •-	
CONT	6908 6910 6912	7202	7412	7501	7712	7801	7803	7805	7807				84	<del> </del>	50.8N	ŗ.	5901	5902	5903	7,904	7000 0000 0000	5907	5908	2909	5912	2009	6003	6004	6009	9009	6007	6000 6010	6102	6104	6107	6201	!
93080	6501 6504	6601	6607	6612	6801	6806	6901 6901	6904	6905 6907	•			93090		30	121	٠.			. ) 🔻		~ ~~	-	-	CU C	uκ	) N.	· <	~	*	$\omega$	$c \sim c$	) r	5707	~ n	0 (7)	3
CONT	7801	7805	7808		· <b>-</b> ·	<u>.</u> . (	s, <b>-</b>	10.8N	ŗ	5904	5905	2900	200gr	5909	5910	1991	6001	6002	6003	4000	6009	6007	6009	6010	6102	6104	6110	6201	6203	6207	6210	7000	6307	6310	6401	6410	) - - )
93070	7202 7203	7501	7712			i d	95080	31	120	5005	5005	5005 5007	5101	5102	5103	5104	7106 7106	5107	5108	5204	2027 5008	5207	5304	5306	5504	7504 5606	5704	5707	5710	5801	5804	7007 7017	5 5 11 11	5812	7901	5902	\ \ \
			· ·		· <b>-</b> ·			· •-·						<b></b> .			· ·	<b>.</b>								·· -		<b>-</b> .	<b>-</b>		<b>-</b>	<b>-</b> -	-		-· <b>-</b>	- · ·	
102	30.8N 14.8W	5909	5911	5912	6002	6003	6004 6005	9009	6007 6008	6010	6102	010	6203	6203	6207	6210	6304	6307	6310	5401	6404	6501	6504	6507	6601	6604	6612	6707	6801	9089	6901	7069	690F	2069	6908 6918	6912	)
93070	31 120	5002 5005	5006	5007	5102	5103	70 40 70 70	5106	5108 5201	5202	5203	7204 7207	1206	5207	5304	5306	5407	5405	5504	2002	7504 7604	5704	5707	5710	5801	7004 F807	5809	5810	5811	5812	5901	7004 1004	1900 1900 1900	5905	5906 1004	5908	
			•			<b></b> .		• •						<b></b> .	<b>-</b>			<b>-</b> .	<b>-</b> ··•		-· <b>-</b> -					<u>-</u>	-	<b>-</b> .				-· <b>-</b>				·· •-	
CON 'T	6104	6201	6207	6210	6307	6310	6401 6404	6410	6501 6504	6507	6601	6604	6610	6612	6707	6801	6901	6902	6904	6907	~ α Ολο Ολο	6910	6912	7202	7203	77.0	7801	7803	7804	7805	7807	808/					
09026	5207 5301 5302	5304 5304	5504	5506 5510	5602	5604	5505 570 <b>4</b>	5707	5710 5801	5804	5807	7804 7810	700	5812	5901	5902	5905	5905	5906	7,000 I	0000	5910	5911	5912	6001	2009 8009	6004	6005	9009	6007	6008	6100	70.10				
<b></b>	. <del></del>		. <del></del> .		. <del></del> .	<b>-</b>			<b></b>		<b>-</b> ·•	<b>-</b> -		· <b>-</b> ·						-· <b>-</b>	<b>-</b> -	٠ 🕳	<b>-</b>			··	٠ ــ.			<b>-</b> ·		- · •-				·· •-	
CON'T	6404 6410 6501	6507	6604	6607	6612	6707	6801 6806	6901	6902 6904	6905	6907	6908	6912	7202	7203	7712	7803	7804	7805	180.7	909				4 0 4			34.3W		5105	5106	50.00 50.00	5202	5203	カンプロ あって A	5206 5206	/ !
93050	5904 5905 5905	5907 5907	5909	5910	5912	6001	5003	6004	6005 6006	2009	6008	6100	6104	6107	6110	6201	6207	6210	6302	0000	070	-			03020	`	_	119		5003	5004	2007 2007	5007	5101	2010	5104	Y
		·	• • •		· <b>-</b>	<b>-</b> ··			<del>_</del> .		<b>-</b> · -	-· <b>-</b> -		<b></b> .					<b>-</b> ·•	<b>-</b> -	·· <u>-</u> -	· <b>-</b> -	<u>۔</u> .				·	<b>-</b>			<b>-</b>	·· -	· <b>-</b> -				
CON'T	7803 7804 7805	888	3			127	Ċ	53.6W	30	2	24	4 <b>&lt;</b>	5 4	5	5	$\frac{4}{6}$	7 =	7	8	2 2	2 5	7.5	9	8	96	56	25	7	8	8	8	2 5	<u> </u>	5812	2 6	88	)
93040 (	6912 7203 7602	3 7 8	3			93050	~	118																										5304			

CALCOFI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

CON 'T	6902 6904 6906 6909 6912 1027	7202 7203 7412 7501	7507 7712 7801 7803 7804	7807 7808 7808	87	15.4N 10.5W 5207	7770 7307 7307 7407 7403 7403 7403 7403	2002 2002 2002 2002 2002 2002 2002 200
97050	710 7801 7801 7080 7090 7090 7000	5000 5000 6001 6001	6501 6601 6607 6612	6806 6901	09026	31 119 5002	2000 2000 2000 2001 2001 2001 2001 2001	5107 5108 5207 5207 5207 5205
<b></b>	. 30.1 30.2 30.1 30.1 30.1 30.1 30.1 50.1							, gant gant gant gant dank dank dank dank
CON'T	7201 7202 7203 7412 7501 7507	7801 7803 7804 7805	7808	90 35.4N 30.1W	5210 5211 5301	2000 2000 2000 2004 2004	2000 2000 2000 2000 2000 2000 2000 200	2000 2000 2000 2000 2001 2001 2001 2001
97040	6507 6601 6601 6607 6707 6806	6908 6908 6904 6907	6910 6910 6912	97050	2002 2005 2006	7007 7101 7102	7010 710 710 710 7110 7110 710 710 710 7	700000 1000000 1000000 1000000 100000000
	, \$60.5 \$50.5 \$50.5 \$50.5 \$50.5 \$50.5 \$50.5						gant gant dank dank dank dank dank	. (m.) (m.) (m.) (m.) (m.) (m.) (m.)
CON'T	7805 7807 7808	55.4N 49.5W	5402 5403 5403 5403 5403	7400 7400 7400 7400 7400 7600 7600	5502 5504 504	5502 5602 5604 5604	7007 7007 7007 7007 7007 7007 7007	0.000 000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.
97035	7803 7804 7804	31	000 000 000 000 000 000 000 000 000 00	2012 2012 2012 2013 2015 2015	5108 5109 1109	5111 5201 5202 5002	00000000000000000000000000000000000000	12000000000000000000000000000000000000
			des des des des des d				er gang dans dans dans dans dans dans dan	s gant gant gant gant gant gant gant
CONT	5904 5907 5910 6001 6002	6004 6005 6707 6801	6902 6904 6906 6907	6912 6912 7201 7412 7501		36 5.4N	69009 69009 69009 70099 70099	7202 7202 7203 7412 7501 7501
97032	5209 5210 5210 5210 5302 5302 5302	2500 2500 2500 2500 200 200 200 200 200	2000 2000 2000 2000 2000 2000	5712 5801 5804 5807 5810 5901		97035	2000 2000 2000 2000 2000 2000 2000 200	5312 6501 6601 6601 6801 6806
								t des ges des des des des des des
48	15.4N 8.8W 5502 5504 5510	5804 5804 5807 5810	7000 7000 7000 7000 7000 7000 7000	6004 7202 7712 7801 7803	7805 7807 7808	9	11.4N 17.0W 5107 5108 5109	5202 5202 5203 5203 5204 5204 5206
97030	71 710 710 7110 7110	72002 - 72002 - 72002 - 7002 - 7002	7200 7200 7200 7200 7200 7001	00000000000000000000000000000000000000	5308 5310 5310	3-70	320 320 3002 5003 5005	2000 2000 2000 2001 2001 2001 2001 2001
<b></b>	., der der 'Ort det der der der de							
CON'T	7203 7807 7808	25 43.5N 27.1W	6904 6905 6908 6909 7201	7202 7203 7205 7205 7803 7804	7805	ω	29.5N 25.3W 7501 7503 7507	
94029	6201 6307 6501 6507	94030 32 117	6004 6010 6101 6301	6310 6401 6404 6712 6804	6806 6902	94032	52 117 6910 6912 7210	
<del>-</del>	er gang gang gang diret diret diret diret ga					.,		
ω	10.8N 55.4W 7202 7804 7807 7808	<u>+</u>	50.8N 35.2W 6907 6910	7202 7804 7807 7808		8 10.8N 54.4W	6910 7804 7807 7808	13 45.5N 23.0W 6104 6107 6110
93110	30 122 5909 6610 6612 6707	93120	29 123 5909 6302	6610 6612 6707 6901 6904		93140 29 124	6707 6901 6904 6907	94029 72 117 6001 6003 6004

8080 21.2N 27.1W 5008 5008 6010 50003 50004 50005 50007 5101 5103 5104 5105 61010 61010 62101 62001 6 7808 CON'T 7000 5011 5101 5102 5103 7805 5003 5004 5005 5006 6912 7201 7202 7412 7412 7412 7403 7403 7404 7404 7405 7407 7407 7407 40000 15.4N 6601 6601 6601 6601 6801 6806 6901 6906 6906 6909 CON 'T 6904 6906 6906 6909 6909 6901 77201 7 2405 2504 2506 120 CONT 77090 77000 77000 77000 77000 77000 77000 77000 77000 77000 77000 77000 77000 77000 77000 77000 77000 50005 50006 50007 5101 5102 5105 5105 5105 5105 5201

OR MORE CRUISES DURING 1950-1978

SIX

CALCOFI STATIONS OCCUPIED ON

7202 7203 7205 7207 7201 7201 7201 7803 7805 7805 7805 7805

CON'T 6407 6501 6601 6601 6602 6902 6904 6906 6906 6907 6910 CON'T 7805 7807 7808 802020 120 7507 7712 7801 7803 1 00090 CRUISES DURING 1950-1978 66909 66 CON 0000 \$\text{Construction for the construction for the co STATIONS OCCUPIED ON 66507 66507 66507 66507 66908 66908 66908 66908 77208 77208 77208 77208 77208 77208 77208 77208 77208 77208 77208 77208 5106 5106 5109 5110 5201 5202 2000 119 CALCOFI CONT 7203 7205 7205 7707 7707 7801 7805 7805 7805 6801 6806 6901 6904 6907 6907 6910 6910 5005 

52	26.9N 44.1W	5910 6001	6004	6612	6707	6901	6902	6904 6906	6907	6910	6912	102/	7203	7207	7507	7801	7803	7805	7807	808/			22		9. 9M	Ċ	5810	5901	5904 5907	5908	5910	6001 6004	7507	
103080	119	5101	5104	5106	5204	5206	5304	5306	5404	5406	5504	0000 1000 1000 1000 1000 1000 1000 100	5704	5710	5804	780.7	700 C	5904	5907	2908			103090	)	626	021	5204	5205	5206	5305	5206	5506 5604	5804	
-		· •- · •-		<b>.</b>		· -·	<b>-</b> ·· <b>-</b>	·· -·				<b>-</b> -	·· •··					<b></b> .	<b>-</b> ···	-· <b>-</b> ·		<b></b> ··		<b>-</b>						. <del>-</del> .				
CONT	7507	7803	7807	ene/			ŭ	10	46.9N	† • •	6004	6612	2100	6801	6901	2069	000 400	6907	6069	6910 6912	7201	7202	7207	7412	7507	7801	7803	7805	7807	2				
103060	6907 6909 6910	6912	7202	7207			040	0/0601	200		5101	2010	5105	5106	5204	5205 506	5304	5305	5206	5504 5506	5604	5704	5707	5801	5804	580 / 5810	5901	5904	5907 5908	5910				
	.,			<b>-</b> .	<b>-</b>	· -·	<b>-</b> · -	<b>-</b> .	<b>-</b> · -		<b>-</b> · .	- · -	<b>.</b>	<b>-</b> ·	<b>-</b> ··•		·· -	· <del>-</del> ·	<b>-</b> ··		<b>.</b> .				<b>-</b> ·•	<b>-</b> -	· <b>-</b>		<b>-</b> · -	٠ ـ.		<b></b>	· • •	
CON 'T	7201 7202 7203	7207	7712	7803	7804	7807	7808			69		N	Z4 · / ×	5804	5807	5810	1000	5903	5904	5905 5906	5907	5908	5910 600 <b>1</b>	6002	6003	6004 7007	9009	2099	6612	6801	9089	6901	6904 6906	: , ,
103050	6612 6707 6801	6806 6901	6902	6904 6906	6907	6979	6912			103060	1	C 4	р 	5101	5103	5104	7105	5204	5205	5304 5305	5306	5401	5402 5403	5404	5405	7406 7500	5504	9055	5512 8602	5604	5701	5704	5710	,
				<b>-</b> -				<b>-</b>		<b>.</b>	<b>-</b> ·•	-· <b>-</b>	<del>-</del>				<u>-</u> -		<b>-</b>		<b>.</b>			. <b>.</b>			· <b>-</b> -			. <b>.</b> .				
CON'T	6901	6906	6069	6910 0120	7201	7203	7207	7503	7507	7712	7801	(300) 100)	7805	7807	7808				95	ý	44.7W		5701	5707	5710	2717 1280	5804	5807	5810	5904	5907	7908 7910	6001	:
103040	5512 5602	5606 5704	5707	5801	5804	5810	5901	5904	5908	6001	6004	2099	5100	6801	9089				103050		117	•	5101	5104	5105	3017 3004	5205	5304	5305	5502	5504	5506 8512	5602 5604	· · · · · · · · · · · · · · · · · · ·
<b></b>			· •	<b>-</b> .		<b>.</b>		<b>-</b> .		·· •·	<b>.</b>	<b>-</b>	-· <b>-</b> ·	<b>-</b> ·			-· <b>-</b>	· <b>-</b>			. <b>-</b> .			· <b>-</b>		<b>-</b> -	<b></b> .	٠.				<b></b>		
CON T	6806 6901 6903	6904 4006	6907	6909 6910	6912	7202	7203	7507	7712	7803	7804	(802	808)				Ca	30	46.9N	•	5307	5308	5310 5310	5401	5402	7405 7405	5405	5406	5407	5410	5412	5502 5504	5506 5510	
103035	5602 5604 5604	5704 5707	5710	5804	5807	5901	5904	5907 5908	5910	6004	2099	6612	6707 6801				103040	0,000	30		5101	5103	ਨ 104 704	5106	5109	5110	5112	5204	5205	5209	5210	5211 5201	5305 5305 5306	/ / / / / / / / / / / / / / / / / / /
~· ~			· •-· •	<b>-</b>		<b>-</b>				-· •	<b>-</b> •		<b>-</b>				-•				. <del>-</del> -						٠ ـ		<b></b> · •	. <b>.</b> .				•
CON 'T	6004 6607 6612																				29	,	56.9N	• Ի	5210	5211	5305	5506	5307	5310	5312	5502 5503	5510	1
103030	5312	5402 5402 5403	5404	5405 5406	5407	5410	5412	5504 5504	5506	5512	5801	7804	580 / 5901	5904	5907	5908 1940	2910				103035		30	-	5103	5104 5105	5106	5109	5110	5112	5204	5205 5205	7208 5208 5209	1
	z >			4 M				· <b>ـ</b> .۔ و		·	•	υ, ••••••	- · ·		~		·· -	. <b>.</b>		<u>-                                    </u>				• •	<u>.</u>	 	· •-·		-· -	·· •·		<u></u>	. <b>-</b>	)
-	1.2N 6.2W	5101	1000	580 650				5	,	45.5W		500	6110	620	620	621	020	640	650	650 720	721				68		24.5W		520	521	530	530 086	5307	) S
100110	122	5002	5004	5007	5009			100120		122		2002	500 <i>5</i>	5005	5007	7101 0101	4017	5404	5804	6008	6101				103030	<b>k</b>	116		5109	5113	5112	5204	5206 5208	i i i

CALCOFI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

CON'T	6902 6904	9069	6069	6910	7201	7202	7207	7412	7507	7801	7803	7805	7807	000			7.7	-	11.5N	o.	200	5804	5807	5810	5901	200	5907	5908	5910	6002	6003	6004	6005	6006 6607	6612	6707	7 - 10
107060	5903	5905 5905	5907	5908 5910	6001	6002	6002	6005	9009	6607	6707	6712	6806	090			107070	)	59	$\infty$	•		-	~	$\alpha$	νс	1 KU	M	10 4	4 4	- 4	S	ഗ	$\sigma$	-	5707	_
			<b></b>					<b></b> .			. <b>-</b> .	<b>-</b>			<b>-</b>						·· <b>-</b>	· <b>-</b> ·			<b>-</b> ··	<b>-</b> -	· <b>-</b> ·	<b>-</b> ·	<b>-</b> •		<b>.</b>						••
CON'T	6712 6806	6901	6904	6906	6069	6910	7201	7202	7203	7207	7510	7712	7801	7804	7805	7807	000			5	2	•	1.3W	,	5606	207	5707	5710	5712	7807	5803	5804	5805	7805 5807	5810	7901	2205
107050	5602 5604	5606	5704	5707	5801	5802	7802 7804	5807	5810	5901 1904	5907	5908	5910	600	6607	6612	10/0			107060	000/01		118					$\sim$	$\alpha$	νĸ	1 K	M	LΩι	ດທ	IN.	5602	ם
	. <b>-</b>		·· ·		<b></b> .			. <b>.</b>			· <b>-</b> ·	<b>-</b>		<del>-</del> -	· <b>-</b>	<b>-</b> ·-	<b>-</b> -	<b></b> .			·· -	·· •·			<b>-</b>				<b></b>						<b>.</b>		
CON T	5910 6001	6004	6612	6707	9089	901	6902 6904	9069	2069	6909 6910	6912	7201	7202	7207	7501	7507	7801	7803	7804	7805	100	000				99		21.6W	•	ড ড	7	4	ུ ५	4 r	/ U	5506	(,)
107040	5210	5304 5304	5306	5307 5308	5310	5312	2002 5504	5506	5510	5512 5602	5604	5606	5701	7077	5710	5801	5804	5810	5901	5904	- 000 - 000 - 000	2900			- 1	060/01		117	i	1017	5104	5105	5204	7207 7207	5304	5305	00/6
			• •		<b></b> .	<b></b> •		· <b>-</b> ·			<b></b> .				· <b>-</b> ·	<b>-··</b>	<b>-</b> -	. <b>-</b>			<b>-</b> -	·· •-·							<b>-</b> ··•				<b></b> ·		<b>-</b> .		••
CONT	5810 5901	5904	5908	5910	6004	6607	5029	6712	9089	6901 6901	6904	9069	6907	6909 6910	6912	7201	7027	7207	7507	7712	1001	7804	7805	7807	7808				70	-	41.8W		(	$\sim$	S	5208	V
107035	5210	5304	5306	5307 5308	5310	5312	7401 7402	5403	5404	5405 5406	5407	5408	5410	74-7 7.00でで	5504	5506	7.7.7 0.1.0	5602	5604	5606	7.04	5710	5801	5804	5807				107040		116		5101	2017 2017	5105	5110	2111
***					<b></b> -			··			· <b>-</b> -				<b>-</b> .		·· •	··			·• •	·• •-·			•	<b>-</b> -	•				·				· <b>-</b> ·		-•
09	27.5N 9.8W	F 00.7	5908	5910	6004	6607	5100	6712	9089	6901 6901	6904	9069	6907	6910	6912	7201	7027	7207	7507	7712	7007	7804	7805	7807	7808				78	_			5112	2204 2004	5206	5208	ンとしか
107032	30	5100	5110	5111	5204	5205	7200 7208	5209	5210	5211	5305	5306	5307	5310	5312	5502	7504	5510	5512	5701	000	5807	5810	5901	5904				107035		116		0	òč	Ô	5110	-
<b>6</b> g			· · ·					. <b>.</b>	<b>-</b> - ·		. 🟎							· <b>-</b> ·			·· -	·· •-·		←.	•	~· <del>-</del>											••
6	9.2N 33.1W	5207	5301	5302	5000				6	σ	12.9W				5303					7	c	52.6W		5207	5301	2006				7	-	9	•	5207	5301	5303	
105050	30	5102	5108	5201	5203				105060		118		5102	5201	5202	5203				105070		118		5102	5201	2000 2007	1607			105080	)	29	119	•	· N	5202	v
	~ ~ ~		• ••	<b></b> -	• •-•	<b></b>		• •			• ••					<b></b>						··						_		-• <del>-</del> -	·						••
I' NO	7808		,	<b>x</b> o	45.2N	<del>.</del>	5207	5301	5302	5303			4	-	39.2N	, '	5108	5201	5202	5203	720	5302	5303					9	ć	77.72	,	5207	5301	5202			
103090 CON'T	7807		1	105032	30	9	71.08	5201	5202	5203			9 2 0 3 0 4	102022	30	9	5002	5003	5004	5005 5005	2000	5007	5011	5102				105040	(	100	,	5102	5108	7201 7200	5203		

126	57.2N 38.7W	5408 5410	5412 5502	5504	5508 5510	5512	5604 4050 4050	5701	5707	5710	7801 801	5803	5804 5807	5810	5905	7983 803	200 400 400	5906	5907	7.00 2.00 2.00 2.00	6001	6002	6003	9009 4008	9009	8009	6010	6105	6107	6111	6203	6207
110060	28	5002 5003	5004 5005	5007	5101 5101 501 501	7103 103 104	5107	5108	5110	5111	5112 5201	5202	5203	5205	5206	5207	000 000 000	5210	5211	2001 2001 2004	5303	5304	5305	7700	5308	5310	5312	5401	5403	5404	5405 5406	5407
CONT	6201 6203 6207	ω ω	ωω	ω α	. W W	ww	<i>y</i> <b>w w</b>	000	υw	Ψ	w u	Φ	w w	w	9	[ · [	- [-	- [	- 1	- [-	- [-	[-	<u>- t</u>	- [	-[-	-	_	•				
110050	5211	5303	5305	5307	5310	5502	5506	5512	5602	5606	5701	1 5707	5710	1 5803	1 5804	5807	7810	1 5904	5907	: 5908 1	6001	1 6004	1 6008	6010	6105	6107	6111					
CON'T	6707 6712 6806	<b>6</b> 6	ω ω	w w	<b>w w</b>	) - [	- [- [	- [- 1			<u>- r</u>	- [-	<u>r</u> r	- [-	[-				108	7	59.2W	\ \	<u>.</u>	2.10	∪ <b>1</b> C	52	52	ν. ν. υ.	57.0	5207	ン (C	52
110040	5904 5907 5908	5910	6004	6010	6105	6111	6203	6211	6302	1 6307	6310	6407	6501	: 6601	1 6612				110050		116		ဥ္ပင္မ	38	38	8	89	55	5	5104	55	0
CONT	7712 7801 7803	7804	7807	)		116	37.2N	•	5307	5310	5312	5402	5403	5405	5406	5407	2408	5412	5502	5504 EE06	5508	5510	5512	5005	7604 5606	5701	5704	5707	5801	5804	5807 5 <b>310</b>	5901
110035	6912 7202 7203	202	212	\ \		110040	29	- 1	$\circ$	$\circ$	$\circ$	$\circ$	5101	~ —	•	<b>-</b> •			<b>.</b>	— ი	u N	$\sim$	$\alpha$	$\sim$ c	VΩ	$\sim$	$\alpha$	$^{\prime\prime}$ $^{\prime\prime}$	110	100	e le	(M)
<b></b>	z z	9 4	· - · -	·	+ r- c	· · - > <	+ [~ 0	00		. <b>.</b>	~··-	· -·		- 10	7	<del>-</del>	N 5	· <b>-</b> ·	0			7	<b>-</b>	- · -	· •	- 0	9.	c	. <del></del>	۰۰۰	~ 6	
104	47.2N 59.8W	വവ	'EV EC	יוטיני	5807	) LL) L	UTUR	ישוי	ی م	ω	<b>6 4</b>	Φ	Φ.	<b>o o</b>	φ	6	υQ	Φ	Ψ	w w	S C	•	Ç (	D U	o w	Φ	w.	D C	Φ	0	D (C	Φ
110035	29	5002	5004	5006	5009	5101	5010	5105	5108 5109	5110	5111	5201	5202	5204	5205	5206	5207 5208	5209	5210	5211	5303	5304	5305	5306	5308	5310	5312	5502	5506	5510	5512 5602	5604
															,																	
49	31.5N 59.5W	5904	5908	6001	7507	78087			r.		51.2N	•	40	54	4	40	<b>₹</b>	1 4	50	20	3.5	7	2	င္ဆ	200	8	8	88	8	5910	38	50
107090	28 119	5304	5506 5604	5606	5804 5804	5810 5810	1060		110033		29 1.15		000	2 =	-	<del></del> (	$\frac{1}{2}$	20	20	200	20	20	2 :	N N	30	30	30	2 K	30	5210	2 2 4	40
F-1	~~~	N	100		•							- CI	~-	<del>-</del> .	<u>-</u> .	-··-	~ 0		<u> </u>	 - C	V K			_ ×	<del>-</del> -		۳		·	•		• •-•
CON 'T	7203 7207 7507	780	780,	780			4	51.5N	ċ	009	600,	661	6707	269	690	069	069	691(	169	720	720	720	750	780	2 2 2 2 2 2 3 2 3 3	780	780					
107070	6901 6902 6904	88	885	200	)		107080	28		5101	5103	5105	5204	5206	5304	5305	5504 5504	5604	2606	5704	5710	5804	5807	5810	7901 7904	5907	5908	5910				

5102 5103 5105 5205 \$2000 \$4000 17.2N 54.0W 22.9N 18.2W 6211 6302 6304 6401 6501 121 6008 6010 6101 6107 6107 6111 5003 5004 5005 5007 5000 6101 6105 6107 6107 6201 6201 6203 6203 6304 6401 6501 5000 5000 5000 5000 5101 5102 5103 6008 57.2N 36.3W CON 'T 64304 5101 5102 5103 119 5003 5004 5005 5007 CON 'T 5202 5204 5204 5205 5207 5301 5304 5404 5403 5908 5009 6009 50005 50005 50005 50005 50005 5 CON'T 6302 6302 6304 66304 66304 66607 66607 66607 66607 66607 6606 6606 6606 6606 6607 

CRUISES DURING 1950-1978 OR MORE SIX NO CALCOFI STATIONS OCCUPIED

117026 CON'T	5301 5710	000 000 040 000	306	308 5	309 5	2 K2 I	702 504 504	506 6 510 6	512 7	609	2	· ·		117030 78	28 47.6N	u ,	202	204	205 205	207	200		302	304	305	307	5309 5701 5310 5708
115035 32	28 55.3V 115 26.7V	1 5108 5009 1 5108 5008								5808 6501 5808 6501				115040 12	28 45.3N	46	יני ט	ט וט	1 5110 5808	יתי	•••	117026 48	ה ה	114 40.1W	c C	203	5204 5209 1 5205 5210 1 5206 5211
113080 CONT	1 7807 7808		113090 10	No. 22 72	9 13.	804	813 913	: 5904 7807 : 5907 7808			115027 14	0	114 55.1W	α	000	5110 5708	112	209			115030 16	29 5.3N 115 7.0W	α C L	100	0 -	- 22	5209 5809 5309 5809
113070 CON!T	5504 6612 5506 6707														115080 38	7 42.	<b>!</b> C	909	604 606	704	710		901	907	908	000	6607 7803 6612 7805
113060 GONT	5306 6003 5412 6004	0000 05004 0006 0006	21.0 2.0 2.0 3.0 5.0 5.0 5.0 5.0 5.0 5.0 5.0 5.0 5.0 5	507 604	606 701	704	707	801	804	810	905	904	905	906	908	910 001	200			113070 62	$\sim$ L	- 2 v	020	200	02	000	5204 5406 5205 5502
113050 CON'T	5304 6001 5305 6002	400 200 201 201	403	404 405	406 502	100	2 5 1 2 5	602 604	606	704 704	710	301 303	304	307 310	901	902 904	907	910			113060 74	28 22.9N 117 16.2W	101 52	102 52	103 52 104 52	105 53	5202 5304 1 5203 5305 1
113040 CON'T	5209 5901 5210 5904	- 00 - 00 - 00 - 00	304	306 306	307 6	310	502 6	504 6 506 6	510 6	602	909	107	707	708 7	710	804	907 7	809 7	5810		113050 76	28 42.9	6 37.	01 52	02 03 50 50	04 52 52	5106 5302 ! 5202 5303 !

CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

-	44.1N 27.7W	5708	5709 5808	5809	5909				•	2	•	47.34	,	5709	8	8	8				12		24.1M	•	9	5708	2	8	င္တ	5									
118025	28 114	5208	5209 5309	5408	5608	5003			0.00	118050	28 8	114	0000	7200 7200	5300	5608	5609				118035		28	115	0003	15209	5209	5408	5411	5608									
	· · ·					-· <u>-</u>							-· <b>-</b>									••••	<b></b> .		- · -		<b>-</b> ·			•					<b>-</b> ··				
CON T	7807 7808				35		10.9W		6712	6902 7002 7004	9069	2069	6909	2.60	7207	7202	7203	7207	7507 7804	7803	7807	7808					တ		47.6M	o.	5907	797 - 0	6001	6004					
117070	7507	508/			117080	7.0	- <del>τ</del>		5506	7504 7506	5704	5707	5710	100 L	5810	5901	5904	5007	5908 1010	6001	6004	6607	6612				117090		5 5 7	118	700 N	्र १८५१	5901	5904					
		-· <del>-</del> -·	<b></b>	<b></b>	<b></b> - •			<b></b>	<b>-</b>		. <b></b> .	<b></b>		·· -		<b></b>		<b>-</b>			• •				- · -		<b>.</b>		<b>-</b> ···		-· <b>-</b>		. <b>.</b>	••	<b>-</b>		·• <b>-</b>		
CONTE	6910 6912	7201	7203	7412	7501	7807	7803	7807	7808			!	7.5	7	30.1W	ł	5801	5804	5807 5810	700.	5904	2007	5908	5910		6507	6612	6712	6901	2069	6904 6904	2009 2009	0009	6910	6912	7201	7007	7207	
117060	6001 5004	6512	6707	6804	6901	7009 9009	6905	4007	60069			1	117070	7	117		5101	5102	5103 5104	7.01 7.05	5106	5202	5203	5204	0707	7007	5303	5304	5305	5306	ひひひ と と と と と と と と に と に と に と に と に ら と に ら と ら と	7 2 C	5512	5602	5604	5606	10 - C	5710	
<b></b>		<b></b> .	<b></b>	· <b>-</b> ·	<b>-</b>				<b>-</b>		<b>.</b>	<b>-</b>	<b>-</b> ·-	<b>-</b> -						. <b>-</b> -	. 🟎	<b>-</b>			- · -		. 🕳	<b>.</b>			<b></b>	·	. 🛶	<b>.</b>		<b>-</b>	<b>-</b>	··	
CON 1.	6902 6904	6906 6907	6909 6910	6912	7201	2027	7207	7507	7510	7801	7807	7808					68		47.6N	•	5406	5505	5504	5506	/ - c - u	7007 7604	5606	5701	5704	5707	5710 5710	- 780 1807	5807	5810	5001	5904	- a 2 C 2 C	540 540	
117050	5904	5906 5907	7908 7910	6001	5005 2005	6003	600F	9009	6607	5512 6707	6712	6804	6901				117060		27	-	5101	5102	5103	5104	7107	7002	5203	5204	5205	5206	5207 5307	7302 7303	5304	5305	5206	5402	7 T U	5405	
				· <del>-</del> ·	<b>-</b> ·•	-· <b>-</b>		. <b>.</b>	<b>-</b> ··		. <b>.</b>			<b>-</b> -		. <b>.</b>					· <b>-</b>				-· <b>-</b>	<b>-</b> -											<b>-</b> -		
CONTE	6612 6707	6712 6804	6901 6902	6904	9069	2.069	6910	6912	7201	72027	7207	7507	7801	7807	7808	) -			1	-	7.64	•		5502	400cc 40cm	77.00 75.10	5602	5604	2606	5704	5707	700 200 200	5803	5804	5807	5810 1001	- 50 H	5903	
117040	5701	5708	5709	5801	5804	7807 808	5809	5810	5901	7904 7907	590R	5909	5910	200	6607	- ) )			117050	2	28	116		$\circ$	$\sim$ $\circ$	$\circ$	$\sim$	$\circ$	$\circ$	$\circ$	5204 5205	$\circ$ $\circ$	$\circ$	$\circ$	$\circ$	$\circ$	$\circ$	5306	
<b></b>							• •	<b></b> .	<b></b> - •	<b>-</b> -	· <b>-</b> ·			·· -		· <b>-</b> ·				-	. <b>.</b>				<b>-</b>		. <b>-</b> .						. <b>-</b>			<b>.</b>	-		
CON 1	6901 6902	6904 6905	6907	6910	6912	7201	7207	7507	7801	7807	7808				88	)	~	35.1W	130B	5300	5510	5401	5402	5403	7404 704	7407 5406	5407	5408	5410	5412	70707 7007	7.7.0.4 5.5.0.4	5510	5512	5602	5604	0000 1000 1000 1000 1000 1000 1000 100	560°	
117035	5808 5809	5810 5901	5904 5907	5908	5909	5910 6004	6004	6607	6612	6707	6804				117040	- >	28	115	-		-	<del></del>	-	<b>-</b> ເ	√ c	uΛ	S	$\sim$	$\sim$	$\sim$	$\sim$	чС	υKU	٣	M	K√ K	NΥ	5307	
		-· -·	<b>-</b>	<b>.</b>	<b></b>	-· <b>-</b>	• •	<b>-</b> .	<b>-</b> ··		<b></b> .	<b>-</b> ···		-· <u>-</u>		. <b>.</b>	<b>-</b>	<b>-</b> ·			. <b>.</b>	<b>-</b> ·	<b>-</b> ••	<b></b>	-· •		<b></b>		<b>-</b> ••		-· <b>-</b> -	·• <del>-</del>		<b>-</b> ·		<b>-</b> · -	<b>.</b> -	<b>.</b>	
CON 'T	6712 6804	6901 6902	6904 6906	6907	6069	6910	7201	7203	7207	7507	7801	7803	7807	909/				72	_	- r - r	•	M	W.	んし	C. 14	UΚ	, <b>K</b>	4	ru.	ທເ	יט יע	$\sim$	$^{\circ}$	[	~	Ç~ 0	υα	5807	
117030	5709	5801 5804	5807 5808	5809	5810	7901	5007	5908	5909	5910 6001	6004	2099	6612	1019				117035		115	-	5003	5004	5005	7000	7007	5011	5101	5102	5103	5104 7107	2000 2000	5203	5208	5209	5211	1000 1000 1000	5303	

60008 60008 60100 7803 7807 7808 7201 7202 7207 7507 120030 CON'T 7807 13.3N 34.3W 2005 5909 5910 6001 6004 7507 7801 7803 114 5709 5710 5804 5808 5808 5809 5810 5901 \times \t CON'T 114 1008 10 6211 6211 6302 6304 66304 66507 66507 6607 6607 6607 6607 6607 16.1N 22.6W 5901 5904 5904 5907 5908 5909 6008 6008 6008 6102 6107 6107 6203 11990 11900 

CALCOFI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

25.0N 40.0W 6203 6211 6304 6401 6501 120 5404 6008 6010 6102 6107 6501 6601 6601 6902 6908 6909 6010 6102 6105 6107 6211 6201 6203 6204 6302 6304 6401 6501 5103 5104 5105 CON'T 6107 6201 6203 6207 6207 6308 6401 5004 5005 5007 5009 7205 7205 7210 7210 7501 7801 7803 CONT 118 6902 6904 6906 6907 6910 6912 7201 33.3N 48.3W CON 'T CON 'T 6712 6712 6804 6804 6906 6907 6907 77203 77 11110 1110 110 110 110 110 110 110 110 110 110 110 110 110 110 110 110 110 110 110 110 1 5910 5010 5010 6000 6902 6906 6906 6912 13.3N 31.0W 116 5504 5502 5602 5604 5606 

6910 6912 7201 7203 7203 7203 7204 7203 7801 7803 7803 65210 652010 652017 652017 652017 653017 653017 653017 653017 653017 653017 653017 43.7N 29.4W 7803 59.0N 25.8W 21420 6001 6004 6607 5707 5710 5804 5810 114 66001 5907 5910 6001 6607 5704 5707 5707 5804 5807 5810 6910 6910 77202 77203 77203 77203 77203 77203 77203 77203 77203 77203 77203 49.0N 5512 5602 1000 5000 5000 5000 5000 5000 5000 5101 5103 6907 6909 6910 6912 77507 77803 7803 7803 59.0N CONT 2000 6401 6401 6501 6507 6612 6712 6904 6904 115 15.0N 59.4W 5003 5005 5005 5009 5011 17.50 17 

CALGOFI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

CALCOFI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

130050 CON'T	5510 6302 5512 6304 5602 6307	604	701	707	801	804 6	807 810 (	901	904	806	016	- 1	008	010	702	107		201	207	211				130060 109		25 29.4N	7 64.	002 51	004 51	005 51	007 51	009 51	01.1	100	5103 5202	
130040 CON'T	6207 6906 6211 6907 6302 6909	304 6 307 6	310 7 401 7	407 7	507 7	601 7	607 7 612 7	707	712 7804 7	902 7	7. 406			•	1,50050 102	5 49.	114 46.2W	В	UL	ıΩ	rU.	וט נ	vπ	ית	וטו	ע ה	U IC	ر م	ı'n	7	וטו	ľΩI	ν.	$\sim$ $\mathbf{r}$	5204 5508	
30040 126	26 9.4N 114 7.9W	5002 5407 5003 5408	004 005	900	600	011	101 102	103	104 105	106	707	0 C	110	111	201	203	204	205	000	208	508	0.70	- C	302	303	504 205	700 700	308	310	512	101	505	403	ל ע כל כ	406	
 E⊣	7807 7808				. 7. V.	<b></b>	306 : 506					<b>-</b> -		<b></b> .	-· -		. <b></b>	-·-	. · ·	· <del>- ·</del>	<b></b> ·	-· -	<b>.</b>	· <del>-</del>	<u></u> .	-·-	<del></del>	. <b>-</b> -	· <del>-</del> ·	<u>-</u> .	<u>.</u>	<b>-</b> -∙ ≀	-· -	·	<del></del> .	
130030 CON	7801 78 7803 78		130035	•	113 48	. '	5002 53		u i u	, 17, 1	7 / 17	7 / L	, E ,	шлі	7 / 17	۱Ц1	ינרו	ΠIR	1 7 E.	, II ,	ורש	TL / E	r · u	, <b>π</b> ,	rv i	υn	ហយ	/ <b>Γ</b> .	·ω	Φ	Γ-					
130030 94	26 29.4N	5108 5809 5109 5810	110	201	203	204	205 206	207	208	210	1 2 2	300	303	304	307 307	307	308	310	401	402	403	404	204 204	407	408	0.4	- t - C - C	504	506	510	512	710		4 C 0	808	
CON'T ! 13	6910 6912 7201	7202	7207 7507 !	7801	7807	1808		•••	p	14		7. (N	. <b></b> .	5901	5907	5910	6001	5004	1000		<b></b> •		رر 	3. T	•			6004	5607	•				·• <del>•</del>	·· •··	
127060 0	5910 6001 6004	607	707	902	906	907	6069	•••	<b>.</b> (m.	127070	T.	116 2	J - -	011	704	710	5801	804	2			•	127080	25 2	7	7	707 707	710	5804	! 5810		<b></b> .				
127050 CON'T	5604 6712 5606 6804 5701 6902	704	71 0 801	804	810	901	904 908	910	001	607	612	2			127060 63	)	26 3.7N	5 46.	ш	, ц ,	ш 🕦	77 / 1	r, n	141	u v	7 / [	. т	νц.	, <b>п</b> ,	п,	<u>π</u> , ι	ц,,	7 / 11	, <b>, u</b>	5206 5908	

70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 701000 70100 70100 70100 70100 70100 70100 70100 70100 7010 133040 CON'T CON'T 7803 7807 7808 7202 7203 7207 55.1N 8.2W 133025 CON'T 29.4N 11.6W 5600 6211 6302 6401 6501 6507 6010 6102 6107 6101 6201 29.4N 6304 6401 6407 6501 6607 7205 7210 CONT 6507 6601 6607 7205 7210 9.4N 56.3W 6211 6302 6304 6401 6501 117 6010 6102 6105 6107 6111 6302 6304 6307 6401 6102 6107 6107 6107 6201 6207 6201 49.4N 40.5W 5707 5710 5804 66008 6610 6105 6107 6107 6201 6203 5003 5003 5004 5005 5102 5103 5103 5103 5201 CON'T 5900 5900 6000 

CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

6107 6107 6107 62001 6200 19.8N 40.4W 2022 5102 5103 5104 5201 5201 5203 MORE CRUISES DURING 1950-1978 64111 65201 65201 65204 65210 65210 65210 65210 65210 65210 65210 65210 65210 65210 65210 77202 77203 7 39.8N 2.5W 5106 5107 5108 5201 2152 5101 5102 5103 CONT 7801 7803 7807 7808 9.8N 5.5W 5904 5904 5904 5908 5910 6004 6501 59.8N 24.5W OR 7201 7202 7203 7203 SIX CALCOFI STATIONS OCCUPIED ON CONT 33.8N 19.8W  $\begin{array}{c} \mathbf{600} \\ \mathbf{600$ 55.1N 55908 55908 66001 66004 66908 66909 66909 72007 35.1N 5907 5908 5910 115 5707 5707 5710 115 

CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

CON'T	5810 5901 5904	5908 6001 6004 7202		18	35.8N 43.0W	5702 5801	5901 5901	5908	6001 6004	1202		15	15.8N	7840	7000 7000 1000 1000 1000	6004 6004	1202			
147025	5502 5512 5602	5604 5702 5704 5801		147030	111	5011	5202	5401	5412	2166	.,	147040	123		50103	5504	5801	tion of space of	gar, ser,	dor: do:
CONT	6001 6004 7202		=	11.2N 57.3W	5904 4908 4908	6004 7202			16	55.8N			7907 7904 1				20	45.8N 24.2W	5109	5202 5301
143050	5810 5901 5904	5908	143060	113	5011	5801 5810	1060	• ••	147020	23	- (	5202	5401	5502	5702		14/025	23	5011	5103 1 5106
CON'T		6,	1.2N 23.4W		7901 7904 7908				;		51.2N 42.2W		7901 1904			t.	<del>.</del>	31.2N 19.8W		5706 5801
143030	7202	143035	112	5011	7202 7202 7202	5502	5602 5604			145040	112	5011	5103	5604	5801		143050	113	5011	: 5103 : 5106
CONTE		_	5.5N 33.9W	6804 7205 7310	0	:	12	9,0	5901	5904 5908	6001 6004	7202		21	11.2N 4.5W	5604 5702	5706 5801	5810 5901	5904 5908	6001 6004
140070	7210	140080	1153	5011	5704		143026	111	5202	5501	5512	5810	<b></b>	143030	24	5011	5109	5501	5412	5512
30	5.5N 41.1W	6001 6111 6201 6203	6211 6302 6304	6401 6401 6407	6507 6804	7205 7205 7210				45.5N 18.8W	5901	5904 5908	6001 6004	7205	0   2)	0	נ	56.44	5508 5704	5908 7205
140050	113	5011 5103 5106	5202 5301	540 5602 5704	5801 5801 5810	5904 5904 5908			140060	114	5011	5103 5106	5109 5602 1304	5801	0.00	140070	23	114	5011	5106
50	35.5N 44.4W	5604 5704 5801	5904 5904	6001 6001 6004	707/	K.	25.5M	× 0 • 0	5908 6001	6004	6201 6203	6211 6302	6304 6401	6501	7202 7202 7205	012/				
140035	24	5011 5103 5106	5202 5301	5412 5502 502 503	2166	140040	22	C11	5011	5106 5109	5202 5301	5502 5508	5512 5602	5704	5705 5801 5810	5901 5904				
23	39.8N 55.8W	6111 6201 6203	6302	6401 6401	6507		56	Ŋ.	25.4W	6111	6203 6211	6302 6304	6401	6507	7202 7205 7210					
137080	115	5704 5707 5710	7000 7000 1000 1000	6001 6001	6102 6105 6107		140030	4	112	5109 5202	5301 5502	5512 5706	5801 5810	5901 5904	5908 6001 6004					

CALCOPI STATIONS OCCUPIED ON SIX OR MORE CRUISES DURING 1950-1978

<del>1</del>	11.9N 3.0W	5702 5708 5708 5801	7202 7202	7210		<del>ر</del> ر	, E	40.2W	5706	7801 7901	6001	7202 7205 7210				13	31.9N 17.2W	5801	5901 6001	7202	2	
157020	110	5011	7.000 2.000 2.000 2.000	5612	<del>-</del> . <del>-</del> -	; ; 1 157030		110	5011	5103	5412	5502 5602 5604	5702	• •	gan e	157040	111	5011	5103	5602	5706	<b></b> .
0 13	1 47.2N 2 15.8W	5801 5 5810 6 5901 6 6001			<u>-</u>	77	2 52.8W	1 5810 5 5901					0 12	2 31.9N	25.		6 5901 9 6001					
153050	11.2	5011 5103 5106 5106	561		1 53060			5011	2200	561			157010	$\sim$		501	5106	541 550	<b></b> •		• •	<b></b>
19	47.2N 24.1W	5706 5708 5801 5810					17	27.2N				5901 5901 5001						7.2N 38.7W		5801 5810		
153020	110	5011 5103 5106 5109	5007 5602 5604	5512 5702 7704	5 / 04		153030	22	- ·	5011	5106	5412 5502 5602	5604 5612	5702			153040	22 111	5011	5103	7507 7604 7612	5702 5704
9	41.5N 51.1W	5109 7205 7210	t	,	1.5N 18.2W	5508	7210			17	ı	9.2W	5708	5810	5901 5908	6001	7202	****				
150070	121	5011 5103 5106	6	0 •	114 2	5011	5106	6016		153016		110	5109	5502	5602 5604	5612 5702	5704					
<b></b>								<b>-</b>	· <del>-</del> · ·	<b></b>					<b></b>		· •-· •-·					
CON 'T	5901 5908	6004 7202 7205 7205		12	21.5N	10.00 R	6001	7202 7202	7210			13	1.5N		5901 5908	6001	7205					
150040	5602 5604 5702	5801 5802 5803 5810		150050	22	7 - CR	5103	5109 5109	5901			150060	22		5011	5106	5401 5508	5810				
ار 	23	0-0rc	o co -	4 (	N IV C	·· <del>-</del> · -		<b>.</b>	· ·	·		- 0 -		- 4	2. 2.	.0	• •-• •		2	z 3	~ ~	- 00
N	11.5N 3.5W	5702 5801 5802 5803	1200g		720	7		-		1.5N 22.2W		5801 5810 5901							0	41.5N 59.6W		5401 5508
150025	111	5011 5103 5106 5109	5202 5301 5401	5412 5502	5512 5512	5604		150030		111	1	5103 5103 5106	5109	5301	5502 5512	5602			150040	111	5011	5106
						<b></b>	•	·· -· -						<b></b> .		~				· • · • · ·		
0	55.8N 58.0W	5904 5908 6001 6004	720%		u	35.3W	5904	600 600 700 700 700 700 700	7202			15	23.5N	<u>+</u>	5801	5901	6001 6004	720%				
147050	22 112	5011 5106 5704 5810	5901	1	147060	113	5011	5106 5704 5810	5901			150019	23	<u>-</u>	5109	5301	5512 5512 5602	5604 5702	ı			

2
1978
┰
ċ
3
1950-19
×
Z
DURING
V.
E3
H
CRUI
$\ddot{z}$
6-7
8
9
R MORE C
Œ
U
OCCUPIED ON SIX OR MORE CRUISES
5
_
Ö
_
IEI
H
Б
Š
8
70
E
2
E
4
SMOILVE
Œ,
50
ĭ
CALCOFI


<u> </u>	, _, _, _, _, _, _, _, _, _, _, _, _,	
	, , , , , , , , , , , , , , , , , , , ,	
21 11.9M 21 11.9M 111 54.2W 5011 5901 5103 6001 5106 7202 5604 7210 5801	20 51.9N 112 31.1W 5011 6001 5602 7202 5604 7205 5901 7210	

TABLE 2. CCFIHYDRO CONTROL CARD SPECIFICATIONS

FIELD 1	FIELD 2	FIELD 3	FIELD 4	FIELD 5	FIELD 6
<title>&lt;/td&gt;&lt;td&gt;DEPTH&lt;/td&gt;&lt;td&gt;TIME&lt;/td&gt;&lt;td&gt;STAT&lt;/td&gt;&lt;td&gt;≺STATIONCODE&gt;&lt;/td&gt;&lt;td&gt;CONCAT&lt;/td&gt;&lt;/tr&gt;&lt;tr&gt;&lt;td&gt;COL 1-24&lt;/td&gt;&lt;td&gt;DISTN&lt;/td&gt;&lt;td&gt;DISTN&lt;/td&gt;&lt;td&gt;STD&lt;/td&gt;&lt;td&gt;&lt;CRUISECODE&gt;&lt;/td&gt;&lt;td&gt;&lt;/td&gt;&lt;/tr&gt;&lt;tr&gt;&lt;td&gt;&lt;/td&gt;&lt;td&gt;DISTP&lt;/td&gt;&lt;td&gt;DISTP&lt;/td&gt;&lt;td&gt;OBS&lt;/td&gt;&lt;td&gt;&lt;STD DEPTH&gt;&lt;/td&gt;&lt;td&gt;PRTREC&lt;/td&gt;&lt;/tr&gt;&lt;tr&gt;&lt;td&gt;&lt;/td&gt;&lt;td&gt;SUBFIL&lt;/td&gt;&lt;td&gt;STATIONS&lt;/td&gt;&lt;td&gt;OBSTD&lt;/td&gt;&lt;td&gt;0&lt;/td&gt;&lt;td&gt;WRTREC&lt;/td&gt;&lt;/tr&gt;&lt;tr&gt;&lt;td&gt;&lt;/td&gt;&lt;td&gt;&lt;/td&gt;&lt;td&gt;CRUISES&lt;/td&gt;&lt;td&gt;ALL&lt;/td&gt;&lt;td&gt;1&lt;/td&gt;&lt;td&gt;PRWREC&lt;/td&gt;&lt;/tr&gt;&lt;tr&gt;&lt;td&gt;&lt;/td&gt;&lt;td&gt;&lt;/td&gt;&lt;td&gt;&lt;/td&gt;&lt;td&gt;&lt;/td&gt;&lt;td&gt;#&lt;/td&gt;&lt;td&gt;#&lt;/td&gt;&lt;/tr&gt;&lt;/tbody&gt;&lt;/table&gt;</title>					

TABLE 3. CCFIHYDRO DATA SELECTION MODES

MODE	CON	TROL CARD SP	ECIFICATIONS		DATA SEL	ECTION CARDS	
	FIELD 2	FIELD 3	FIELD 4	FIELD 5	1st Set	2ND SET	3RD SET
1	DEPTH	DISTN OR STATIONS	STAT	CRUISECODE OR O OR 1	0	STATIONS	CRUISES (IF FIELD 5 = 1)
2	DEPTH	TIME OR CRUISES	STAT	STATIONCODE OR O	0	CRUISES	NONE
3	DEPTH	DISTP	STAT	CRUISECODE OR O OR 1	0	STATIONS	CRUISES (IF FIELD 5 = 1)
4	DISTN	TIME OR CRUISES	STAT	0	STATIONS	CRUISES	NONE
5	DISTP	DISTN	STAT	CRUISECODE OR O OR 1	0	STATIONS	CRUISES (IF FIELD 5 =1)
6	DISTP	TIME OR CRUISES	STAT	0	STATIONS	CRUISES	NONE
7	DEPTH	DISTN OR STATIONS	STD	CRUISECODE OR O OR 1	DEPTHS	STATIONS	CRUISES (IF FIELD 5 = 1)
8	DEPTH	TIME OR CRUISES	STD	STATIONCODE OR O	DEPTHS	STATIONS	CRUISES (IF
9	DEPTH	DISTP	STD	CRUISECODE OR O OR 1	DEPTHS	STATIONS	CRUISES (IF FIELD 5 = 1)
10	DISTM	TIME OR CRUISES	STD	STD DEPTH	STATIONS	CRUISES	NONE
11	DISTP	DISTN	STD	CRUISECODE OR O OR 1	STD DEPTH	STATIONS	CRUISES (IF FIELD 5 = 1)
12	DISTP	TIME OR CRUISES	STD	STD DEPTH	STATIONS	CRUISES	NONE
13	DEPTH	DISTN OR STATIONS	OBS OR OBSTD	CRUISECODE OR O OR 1	SEQ NOS	STATIONS	CRUISES (IF FIELD 5 = 1)
14	DEPTH	TIME OR CRUISES	OBS OR OBSTD	STATIONCODE OR O	SEQ NOS	CRUISES	NONE
15	DEPTH	DISTP	OBS OR OBSTD	CRUISECODE OR O OR 1	SEQ NOS	STATIONS	CRUISES (IF FIELD 5 = 1)
16	<del>-</del>	<b>-</b> .	-	NOT USE	D -	-	-
17	SUBFIL	STATIONS	ALL	#	STATIONS	NONE	NONE
18	SUBFIL	CRUISES	ALL	#	CRUISES	NONE	NONE



## **RECENT TECHNICAL MEMORANDUMS**

Copies of this and other NOAA Technical Memorandums are available from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22167. Paper copies vary in price. Microfiche copies cost \$3.50. Recent issues of NOAA Technical Memorandums from the NMFS Southwest Fisheries Center are listed below:

- NOAA-TM-NMFS-SWFC- 14

  Histological gonad analyses of late summer—early winter collections of bigeye tuna, *Thunnus obesus*, and yellowfin tuna, *Thunnus albacares*, from the Northwest Atlantic and the Gulf of Mexico.

  S. R. GOLDBERG and H. HERRING-DYAL (June, 1981)
  - Status reports on world tuna and billfish stocks.STAFF, SWFC (July, 1981)
  - An evaluation of tagging, marking, and tattooing techniques for small delphinids.
     M. J. WHITE, JR., J. G. JENNINGS, W. F. GANDY and L. H. CORNELL
    - (November 1981)
  - Local stability in maximum net productivity levels for a simple model of porpoise population sizes.
     T. POLACHECK

(April 1981)

- 18 Environmental data contouring program EDMAP2. L. EBER (April 1982)
- The relationship between changes in gross reproductive rate and the current rate of increase for some simple age structured models.
  T. POLACHECK
  (May 1982)
- Testing methods of estimating range and bearing to cetaceans aboard the R/V D. S. Jordan.
   T. D. SMITH (1982)
- 21 "An annotated bibliography of the ecology of co-occurring tunas (Katsuwonus pelamis, Thunnus albacares) and dolphins (Stenella attenuata, Stenella longirostris and Delphinus delphis in the eastern tropical Pacific"

S. D. HAWES (November 1982)

- 22 Structured flotsam as fish aggregating devices. R. S. SHOMURA and W. M. MATSUMOTO (November 1982)
- Abundance estimation of dolphin stocks involved in the eastern tropical Pacific yellowfin tuna fishery determined from aerial and ship surveys to 1979.
   R. S. HOLT and J. E. POWERS (November 1982)